

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/EP04/014170

International filing date: 13 December 2004 (13.12.2004)

Document type: Certified copy of priority document

Document details: Country/Office: EP  
Number: 04027213.0  
Filing date: 16 November 2004 (16.11.2004)

Date of receipt at the International Bureau: 28 January 2005 (28.01.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



**Europäisches  
Patentamt**

**European  
Patent Office**

**Office européen  
des brevets**

**Bescheinigung**

**Certificate**

**Attestation**

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

**Patentanmeldung Nr. Patent application No. Demande de brevet n°**


04027213.0

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

**R C van Dijk**





Anmeldung Nr:  
Application no.: 04027213.0  
Demande no:

Anmeldetag:  
Date of filing: 16.11.04  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Epigenomics AG  
Kastanienallee 24  
10435 Berlin  
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se referer à la description.)

PIXT2 - a marker to predict survival of patients diagnosed with breast cell  
proliferative disease

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)  
revendiquée(s)

Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

G01N33/48

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LU MC  
NL PL PT RO SE SI SK TR LI





16. Nov. 2004

**PITX2 - a marker to predict survival of patients diagnosed with breast cell proliferative disease**

The present invention relates to methods for predicting the survival of a human being diagnosed with a cell proliferative disorder of the breast tissues, characterised by a step of determining the expression level of PITX2 or the genetic or the epigenetic modifications of the genomic DNA associated with the gene PITX2. The invention also relates to sequences, oligonucleotides and antibodies which can be used within the described methods.

**Field of the Invention****BREAST CANCER SURVIVAL**

In European and American women breast cancer is the most frequently diagnosed cancer and the second leading cause of cancer death. In women aged 40-5, breast cancer is the leading cause of death (Greenlee *et al.*, 2000). In 2002 there were 204,000 new cases of breast cancer in the US and a comparable number in Europe.

Breast cancer is defined as the uncontrolled proliferation of cells within breast tissues. Breasts are comprised of 15 to 20 lobes joined together by ducts. Cancer arises most commonly in the duct, but is also found in the lobes with the rarest type of cancer termed inflammatory breast cancer. It will be appreciated by those skilled in the art that there exists a continuing need to improve methods of early detection, classification and treatment of breast cancers. In contrast to the detection of some other common cancers such as cervical and dermal there are inherent difficulties in classifying and detecting breast cancers.

The first step of any treatment is the assessment of the patient's condition comparative to defined classifications of the disease. However the value of such a system is inherently dependent upon the quality of the classification. Breast cancers are staged according to their size, location and occurrence of metastasis. Methods of treatment include the use of surgery, radiation therapy, chemotherapy and endocrine therapy, which are also used as adjuvant therapies to surgery. In general more aggressive diseases should be treated with more aggressive therapies.

Although the vast majority of early cancers are operable, i.e. the tumor can be completely removed by surgery, about one third of the patients with lymph-node negative diseases and about 50-60% of patients with node-positive disease will develop metastases during follow-up.

Based on this observation, systemic adjuvant treatment has been introduced for both node-positive and node-negative breast cancers. Systemic adjuvant therapy is administered after surgical removal of the tumour, and has been shown to reduce the risk of recurrence significantly. Several types of adjuvant treatment are available: endocrine treatment, also called hormone treatment (for hormone receptor positive tumours), different chemotherapy regimens, and antibody treatments based on novel agents like Herceptin (an antibody to an epidermal growth factor receptor).

The growth of the majority of breast cancers (app. 70-80%) is dependent on the presence of estrogen. Therefore, one important target for adjuvant therapy is the removal of estrogen (e.g. by ovarian ablation) or the blocking of its synthesis or the blocking of its actions on the tumour cells either by blocking the receptor with competing substances (e.g. Tamoxifen) or by inhibiting the conversion of androgen into estrogen (e.g. aromatase inhibitors). This type of treatment is called "endocrine treatment". Endocrine treatment is thought to be efficient only in tumours that express hormone receptors (the estrogen receptor (ER) and/or the progesterone receptor (PR)). Currently, the vast majority of women with hormone receptor positive breast cancer receive some form of endocrine treatment, independent of their nodal status. The most frequently used drug in this scenario is Tamoxifen.

However, even in hormone receptor positive patients, not all patients benefit from endocrine treatment. Adjuvant endocrine therapy reduces mortality rates by 22% while response rates to endocrine treatment in the metastatic (advanced) setting are 50 to 60%.

Since Tamoxifen has relatively few side effects, treatment may be justified even for patients with low likelihood of benefit. However, these patients may require additional, more aggressive adjuvant treatment. Even in earliest and least aggressive tumours, such as node-negative, hormone receptor positive tumours, about 21% of patients relapse within 10 years after initial diagnosis if they receive Tamoxifen monotherapy only, as adjuvant treatment (Lancet, 1998

May 16;351(9114):1451-67. Tamoxifen for early breast cancer: an overview of the randomised trials. Early Breast Cancer Trialists' Collaborative Group.). Similarly, some patients with hormone receptor negative disease may be treated sufficiently with surgery and potentially radiotherapy alone, whereas others may require additional chemotherapy.

Several cytotoxic regimens have shown to be effective in reducing the risk of relapse in breast cancer (Mansour *et al.*, Survival advantage of adjuvant chemotherapy in high-risk node-negative breast cancer: ten-year analysis--an intergroup study. J Clin Oncol. 1998 Nov;16(11):3486-92.). According to current treatment guidelines, most node-positive patients receive adjuvant chemotherapy both in the US and Europe, since the risk of relapse is considerable. Nevertheless, not all patients do relapse, and there is a proportion of patients who would never have relapsed even without chemotherapy, but who nevertheless receive chemotherapy due to the currently used criteria. In hormone receptor positive patients, chemotherapy is usually given before endocrine treatment, whereas hormone receptor negative patients receive only chemotherapy.

The situation for node-negative patients is particularly complex. In the US, cytotoxic chemotherapy is recommended for node-negative patients, if the tumour is larger than 1 cm. In Europe, chemotherapy is considered for the node-negative cases if one or more risk factors such as tumour size larger than 2 cm, negative hormone receptor status, or tumour grading of three or age <35 is present. In general, there is a tendency to select premenopausal women for additional chemotherapy whereas for postmenopausal women, chemotherapy is often omitted. Compared to endocrine treatment, in particular Tamoxifen or aromatase inhibitors, chemotherapy is highly toxic, with short-term side effects such as nausea, vomiting, bone marrow depression, and long-term effects such as cardiotoxicity and an increased risk for secondary cancers.

It is currently not clear which breast cancer patients should be selected for more aggressive therapy and which would do well without additional aggressive treatment, and clinicians agree that there is a large need for proper selection of patients. The difficulty of selecting the right patients for adjuvant treatment and selecting the right adjuvant treatment, and the lack of suitable criteria is also reflected by a recent study which showed that chemotherapy is used much less frequently than recommended, based on data from the New Mexico Tumor registry

(Du *et al.*, 2003). This study provided substantial evidence that there is a need for better selection of patients for chemotherapy or other, more aggressive forms of breast cancer therapy.

PITX2 (also known as PTX2, RS, RGS, ARP1, Brx1, IDG2, IGDS, IHG2, RIEG, IGDS2, IRID2, Otlx2, RIEG1, MGC20144) is known to belong to the PTX subfamily of PTX1, PTX2, and PTX3 genes which define a novel family of transcription factors, within the paired-like class of homeodomain factors. The gene PITX2 (according to NM\_153426) encodes the paired-like homeodomain transcription factor 2, which is known to be expressed during development of anterior structures such as the eye, teeth, and anterior pituitary.

Toyota *et al.*, (2001) (Blood 97: p 2823-9.) found hypermethylation of the PITX2 gene in a large proportion of acute myeloid leukemias. Furthermore, in this study hypermethylation of PITX2 is positively correlated to methylation of the ER gene and to a reduced expression level. Means to analyse the methylation pattern of the PITX2 gene have been described in a number of patent applications, too (WO 02/077272 is related to the use of methylation markers to differentiate between AML and ALL, WO 01/19845 is related to several differentially methylated sequences useful for diagnosis of several cell proliferative disorders, WO 02/00927 and WO 01/092565 are related to the use of methylation markers to diagnose diseases associated with development genes or associated with DNA transcription, respectively. Loss of heterozygosity (hereinafter also referred to as 'LOH') of chromosome 4 is a known characteristic of many tumour types. Shivapurkar *et al.* [*Cancer Research* 59, 3576-3580, August 1, 1999] have observed loss of heterozygosity at multiple regions of chromosome 4 in breast cancer samples and cell lines. Deletions at 4q25-26 were present in 67% of analysed samples. However the analysed region (between markers D4S1586 and D4S175) does not map to the PITX2 gene, and no inference concerning PITX2 expression was made. Furthermore, the investigation as carried out does not indicate the suitability of any genes or loci of the region for a prognostic use.



Although the methylation of PITX2 has been associated with development, transcription and disease such as cancer, it has no heretofore recognised role in the outcome prediction of breast cancer patients or responsiveness to endocrine treatment.

## EXPRESSION ANALYSIS

The expression of a gene, or rather the protein encoded by the gene, can be studied on four different levels: firstly, protein expression levels can be determined directly, secondly, mRNA transcription levels can be determined, thirdly, epigenetic modifications, such as gene's DNA methylation profile or the gene's histone profile; can be analysed, as methylation is often correlated with inhibited protein expression, and fourth, the gene itself may be analysed for genetic modifications such as mutations, deletions, polymorphisms etc. influencing the expression of the gene product.

The levels of observation that have been studied by the methodological developments of recent years in molecular biology, are the genes themselves, the transcription of these genes into RNA, and the translation into the resulting proteins. However how the activation and inhibition of specific genes, in specific cells and tissues, at specific time points in the course of development of an individual are controlled, is correlatable to the degree and character of the methylation of the genes or respectively the genome. In this respect, pathogenic conditions may manifest themselves in a changed methylation pattern of individual genes or of the genome.

The four terms that apply to the fields of overall genome-wide analysis of all these biological processes are called: Proteomics, Transcriptomics, Epigenomics (or Methylomics) and Genomics. Methods and techniques that can be used for studying expression or studying the modifications responsible for expression on all of these levels are well described in the literature and therefore known to a person skilled in the art. They are described in text books of molecular biology and in a large number of scientific journals.

How to analyse the protein expression of a single gene is prior art. It usually requires an antibody specific for the gene product of interest. Appropriate technologies would be ELISA or Immunohistochemistry.

The analysis of the level of mRNA also has been described sufficiently. These days the gold standard is the reverse transcriptase PCR.

To avoid duplication a more detailed description of the prior art relating to existing and well known technologies is given within the description of the invention, as it is part of the invention.

US patent application 2003/0198970 by Gareth Roberts lists some of the technologies and methods on how to determine a person's "genetic make up", i.e. the genetic modifications, such as deletions, polymorphisms, mutations etc. that may vary between individuals and describes the potential role of this genetic sequence information in the individual's variability in disease, response to therapy and prognosis. Epigenetic differences however are not mentioned. The gene PITX2 is listed within this application as one gene name out of a long and comprehensive list of about 2.500 other gene names, suggesting its expression could play a role in some kind of treatment response. However, this is simply an assumption based on speculation only, as no experiments are disclosed, which demonstrate any kind of relation between genetic modifications of PITX2 and an individual's variation in treatment response.

A less established area in this context is the field of epigenomics or epigenetics, i.e. the field concerned with analysis of DNA methylation patterns.

Methylation of DNA can play an important role in the control of gene expression in mammalian cells. DNA methyltransferases are involved in DNA methylation and catalyse the transfer of a methyl group from S-adenosylmethionine to cytosine residues to form 5-methylcytosine, a modified base that is found mostly at CpG sites in the genome. The presence of methylated CpG islands in the promoter region of genes can suppress their expression. This process may be due to the presence of 5-methylcytosine, which apparently interferes with the binding of transcription factors or other DNA-binding proteins to block transcription. In different types of tumours, aberrant or accidental methylation of CpG islands in the promoter region has been observed for many cancer-related genes, resulting in the silencing of their expression. Such genes include tumour suppresser genes, genes that suppress metastasis and angiogenesis, and genes that repair DNA (Momparker and Bovenzi (2000) J. Cell Physiol. 183:145-54).

In addition it has been described that DNA methylation may also play a role in the field of pharmacogenetics. A similar approach on how to apply information about genetic modifications of the genome to the analysis of individual responses to treatment as was for example described by Gareth Roberts in US application 2003/0198970 was already subject of the application WO 02/037398, tailored to the application of information about *epigenetic* modifications of the genome, based on DNA methylation analysis, to guide treatment selection and to study individual's treatment responses.

An example for the applicability of this idea was given by Esteller et al. (Esteller et al. (2000) N Engl J Med. 2000 Nov 9;343(19):1350-4.), who demonstrated that methylation of the MGMT promoter in gliomas is a useful predictor of the responsiveness of the tumours to alkylating agents. More recently, Frühwald has summarised a series of studies demonstrating that DNA methylation is associated with the aggressiveness of different cancers (Frühwald MC. DNA methylation patterns in cancer: novel prognostic indicators? Am J Pharmacogenomics. 2003;3(4):245-60).

An example for the potential of analysis of epigenetic modifications, such as DNA methylation analysis, for the prediction of treatment response - related to breast cancer- was presented as a poster by Martens et al. at the San Antonio Breast Cancer Symposium, San Antonio, TX, December, 3-6, 2003. Breast cancer patients which have had their tumours removed by surgery and developed metastases at some point after the removal, were treated with Tamoxifen, an endocrine treatment drug. The primary tumour samples were analysed for aberrant methylation patterns. The patients were then divided into two sub classes according to their objective tumour response: patients with progressive disease (which could be described as increasing metastasis size) and patients with complete or partial remission of the relapsed tumour (which could be described as decreasing metastasis size). It turned out, that those patients which had a tumour removed and experienced a remission (decrease in size) of the metastasis under endocrine treatment, had suffered from a tumour which showed a distinct pattern of DNA methylation at specific CpG sites, whereas patients which show progressive disease (did not experience a decrease but an increase in size of their metastases), under endocrine treatment, suffered from a tumour which did not show this distinct pattern of DNA methylation (but a different pattern) at these CpG sites. This is a clear indication, that the methylation pattern described in that study can serve as a **predictive** treatment response tool for an endocrine treatment, like tamoxifen. The results of this study, i.e. predictive biomarkers and assays



therefore, are subject of patent application WO 04/035803, published at April 29, 2004: Method and nucleic acid for the improved treatment of breast cell proliferative disorders. Predictive markers as described above will also be called 'metastatic' markers in the context of this application. PITX2 is also listed as a predictive marker in said application.

Currently several predictive markers are under evaluation. As up to now most patients have received Tamoxifen as endocrine treatment most of the markers have been shown to be associated with response or resistance to tamoxifen. However, it is generally assumed that there is a large overlap between responders to one or the other endocrine treatment. In fact, ER and PR expression are used to select patients for any endocrine treatment. Among the markers which have been associated with tamoxifen response is bcl-2. High bcl-2 expression levels showed promising correlation to tamoxifen therapy response in patients with metastatic disease and prolonged survival and added valuable information to an ER negative patient subgroup (J Clin Oncology, 1997, 15 5: 1916-1922; Endocrine, 2000, 13(1):1-10). There is conflicting evidence regarding the independent predictive value of c-erbB2 (Her2/neu) overexpression in patients with advanced breast cancer that require further evaluation and verification (British J of Cancer, 1999, 79 (7/8):1220-1226; J Natl Cancer Inst, 1998, 90 (21): 1601-1608).

Other predictive markers include SRC-1 (steroid receptor coactivator-1), CGA mRNA over expression, cell kinetics and S phase fraction assays (Breast Cancer Res and Treat, 1998, 48:87-92; Oncogene, 2001, 20:6955-6959). Recently, uPA (Urokinase-type plasminogen activator) and PAI-1 (Plasminogen activator inhibitor type 1) together showed to be useful to define a subgroup of patients who have worse prognosis and who would benefit from adjuvant systemic therapy (J Clinical Oncology, 2002, 20 n° 4). However, all of these markers need further evaluations in prospective trials as none of them is yet a validated marker of response.

Also recently published was a study related to the **prognostic** power of methylation analysis in breast cancer patients. Müller et al. (Muller HM, Widschwendter A, Fiegl H, Ivarsson L, Goebel G, Perkmann E, Marth C, Widschwendter M. (2003) DNA methylation in serum of breast cancer patients: an independent prognostic marker. Cancer Res. 2003 Nov 15; 63(22): 7641-5.) reported about a set of genes, which can be used as biomarkers in patient pre-therapeutic sera for the prognosis of breast cancer. Specific aberrant methylation patterns of

two genes found in DNA from pretreatment serum of cancer patients indicated whether their prognosis was good or bad. The DNA analysed was not tumour DNA but serum DNA. Most likely the presence of a tumour-specific pattern indicates that tumour derived DNA is present, however, the absence of a specific methylation pattern, may be due to a tumour which does not show this methylation pattern, or a tumour which does not shed sufficient DNA into the blood stream. Good or bad prognosis was defined as long or short "overall survival" after surgery, without adjuvant treatment. This result therefore relates to untreated patients, only.

These 'prognostic' markers are able to answer the question whether or not a breast cancer patient should get an aggressive adjuvant treatment like chemotherapy after removal of the tumour to avoid recurrence of cancer, i.e. occurrence of metastases.

However, none of these study results and none of these markers is able to answer the specific question raised above, whether or not a breast cancer patient should get adjuvant chemotherapy after removal of the tumour to avoid recurrence of cancer, i.e. occurrence of metastases in addition or alternative to endocrine treatment (with a drug like tamoxifen, or aromatase inhibitors).

A marker for a bad prognosis for cancer patients (without treatment), might not be applicable to a patient under adjuvant treatment with a drug like tamoxifen. Therefore the test would not be able to help deciding, whether chemotherapy, including all its side affects and inherent risks, is necessary or whether endocrine treatment is sufficient, because an endocrine treatment might change the prognosis from "bad" to "good".

The predictive 'metastatic' marker set described above, would be able to identify amongst all patients which relapsed (developed metastases after surgery) those patients, which do not respond to endocrine treatment (by partial or complete remission of relapsed tumour). These markers however, cannot be applied to answer the question whether metastases will occur at all (after surgery of the primary tumour under endocrine treatment), and consequently whether it is advised to give adjuvant chemotherapy to avoid recurrence of cancer (i.e. relapse or occurrence of metastases).

In one aspect the present invention provides a marker, PITX2 (which shall be recognised as the gene encoding for the protein PITX2; according to NM\_153426), that can be used to an-

answer that question and help guiding the decision whether or not an adjuvant chemotoxic therapy shall be subscribed in addition or instead of treatment with endocrines, like tamoxifen. A marker able to answer this question will also be called 'adjuvant' marker, in the context of this application.

The herein described invention provides a novel breast cell proliferative disorder prognostic biomarker.

It is herein disclosed that aberrant expression of the gene PITX2 is correlated to prognosis of breast cell proliferative disorder patients, in particular breast carcinoma.

In particular this marker provides a novel means for the characterisation of breast carcinomas. Aberrant expression of the gene PITX2 is indicative of the survival of a breast carcinoma patient treated with one or more treatments which target the estrogen receptor, synthesis or conversion pathways or are otherwise involved in estrogen metabolism, production or secretion. The herein described invention is particularly useful for the differentiation of individuals who may be appropriately treated with one or more treatments which target the estrogen receptor pathway or are involved in estrogen metabolism, production or secretion from those individuals who would be optimally treated with other treatments in addition to said treatment. Preferred 'other treatments' include but are not limited to chemotherapy or radiotherapy.

As used herein the term expression shall be taken to mean the transcription and translation of a gene. The level of expression of a gene may be determined by the analysis of any factors associated with or indicative of the level of transcription and translation of a gene including but not limited to methylation analysis, loss of heterozygosity (hereinafter also referred to as LOH), RNA expression levels and protein expression levels.

Furthermore the activity of the transcribed gene may be affected by genetic variations such as but not limited genetic mutations (including but not limited to SNPs, point mutations, deletions, insertions, repeat length, rearrangements and other polymorphisms).

In addition study results presented by Paik et al. at the San Antonio Breast Cancer Symposium, San Antonio, TX, December, 3-6, 2003 provide an answer to this question, by analysing the mRNA expression pattern of 16 genes plus 5 controls with RT-PCR.



The provided invention however has the advantage that looking at only one gene or a small selection of three to five genes will give sufficient information for a validated prognosis.

For demonstration : The 'metastatic' test (use of a 'metastatic' marker) tells a patient whether she is unlikely to respond to endocrine treatment when she develops metastases. But she does not know how high the likelihood is, that she will experience a relapse at all. The 'prognostic' test (use of a 'prognostic' marker) tells a patient whether she will have a good or bad prognosis without any treatment. Even with a "bad prognosis" endocrine treatment might be enough though. The prognostic markers are not necessarily able to predict the outcome under endocrine treatment. The 'adjuvant test' (use of an 'adjuvant' marker) tells her whether she will or will not develop recurrence, without chemotherapy, even when treated with the standard -low side effected- endocrine treatment.

This invention relates to the use of PITX2, as an 'adjuvant marker', which also serves as a 'prognostic marker', especially in hormone receptor negative women, which would not get any endocrine treatment at all.

5-methylcytosine is the most frequent covalent base modification in the DNA of eukaryotic cells. It plays a role, for example, in the regulation of the transcription, in genetic imprinting, and in tumorigenesis. Therefore, the identification of 5-methylcytosine as a component of genetic information is of considerable interest. However, 5-methylcytosine positions cannot be identified by sequencing since 5-methylcytosine has the same base pairing behaviour as cytosine. Moreover, the epigenetic information carried by 5-methylcytosine is completely lost during PCR amplification.

A relatively new and currently the most frequently used method for analysing DNA for 5-methylcytosine is based upon the specific reaction of bisulfite with cytosine which, upon subsequent alkaline hydrolysis, is converted to uracil which corresponds to thymidine in its base pairing behaviour. However, 5-methylcytosine remains unmodified under these conditions. Consequently, the original DNA is converted in such a manner that methylcytosine, which originally could not be distinguished from cytosine by its hybridisation behaviour, can now be detected as the only remaining cytosine using "normal" molecular biological techniques, for example, by amplification and hybridisation or sequencing. All of these techniques are based

on base pairing which can now be fully exploited. In terms of sensitivity, the prior art is defined by a method which encloses the DNA to be analysed in an agarose matrix, thus preventing the diffusion and renaturation of the DNA (bisulfite only reacts with single-stranded DNA), and which replaces all precipitation and purification steps with fast dialysis (Olek A, Oswald J, Walter J. A modified and improved method for bisulphite based cytosine methylation analysis. *Nucleic Acids Res.* 1996 Dec 15;24(24):5064-6). Using this method, it is possible to analyse individual cells, which illustrates the potential of the method. However, currently only individual regions of a length of up to approximately 3000 base pairs are analysed, a global analysis of cells for thousands of possible methylation events is not possible. However, this method cannot reliably analyse very small fragments from small sample quantities either. These are lost through the matrix in spite of the diffusion protection.

An overview of the further known methods of detecting 5-methylcytosine may be gathered from the following review article: Rein, T., DePamphilis, M. L., Zorbas, H., *Nucleic Acids Res.* 1998, 26, 2255.

To date, barring few exceptions (e.g., Zeschnigk M, Lich C, Buiting K, Doerfler W, Horsthemke B. A single-tube PCR test for the diagnosis of Angelman and Prader-Willi syndrome based on allelic methylation differences at the SNRPN locus. *Eur J Hum Genet.* 1997 Mar-Apr;5(2):94-8) the bisulfite technique is only used in research. Always, however, short, specific fragments of a known gene are amplified subsequent to a bisulfite treatment and either completely sequenced (Olek A, Walter J. The pre-implantation ontogeny of the H19 methylation imprint. *Nat Genet.* 1997 Nov;17(3):275-6) or individual cytosine positions are detected by a primer extension reaction (Gonzalzo ML, Jones PA. Rapid quantitation of methylation differences at specific sites using methylation-sensitive single nucleotide primer extension (Ms-SNuPE). *Nucleic Acids Res.* 1997 Jun 15;25(12):2529-31, WO 95/00669) or by enzymatic digestion (Xiong Z, Laird PW. COBRA: a sensitive and quantitative DNA methylation assay. *Nucleic Acids Res.* 1997 Jun 15;25(12):2532-4). In addition, detection by hybridisation has also been described (Olek et al., WO 99/28498).

Further publications dealing with the use of the bisulfite technique for methylation detection in individual genes are: Grigg G, Clark S. Sequencing 5-methylcytosine residues in genomic DNA. *Bioessays.* 1994 Jun;16(6):431-6, 431; Zeschnigk M, Schmitz B, Dittrich B, Buiting K, Horsthemke B, Doerfler W. Imprinted segments in the human genome: different DNA meth-

ylation patterns in the Prader-Willi/Angelman syndrome region as determined by the genomic sequencing method. *Hum Mol Genet.* 1997 Mar;6(3):387-95; Feil R, Charlton J, Bird AP, Walter J, Reik W. Methylation analysis on individual chromosomes: improved protocol for bisulphite genomic sequencing. *Nucleic Acids Res.* 1994 Feb 25;22(4):695-6; Martin V, Ribieras S, Song-Wang X, Rio MC, Dante R. Genomic sequencing indicates a correlation between DNA hypomethylation in the 5' region of the pS2 gene and its expression in human breast cancer cell lines. *Gene.* 1995 May 19;157(1-2):261-4; WO 97/46705 and WO 95/15373.

An overview of the Prior Art in oligomer array manufacturing can be gathered from a special edition of *Nature Genetics* (*Nature Genetics Supplement*, Volume 21, January 1999), published in January 1999, and from the literature cited therein.

Fluorescently labelled probes are often used for the scanning of immobilised DNA arrays. The simple attachment of Cy3 and Cy5 dyes to the 5'-OH of the specific probe are particularly suitable for fluorescence labels. The detection of the fluorescence of the hybridised probes may be carried out, for example via a confocal microscope. Cy3 and Cy5 dyes, besides many others, are commercially available.

Matrix Assisted Laser Desorption Ionization Mass Spectrometry (MALDI-TOF) is a very efficient development for the analysis of biomolecules (Karas M, Hillenkamp F. Laser desorption ionization of proteins with molecular masses exceeding 10,000 daltons. *Anal Chem.* 1988 Oct 15;60(20):2299-301). An analyte is embedded in a light-absorbing matrix. The matrix is evaporated by a short laser pulse thus transporting the analyte molecule into the vapour phase in an unfragmented manner. The analyte is ionised by collisions with matrix molecules. An applied voltage accelerates the ions into a field-free flight tube. Due to their different masses, the ions are accelerated at different rates. Smaller ions reach the detector sooner than bigger ones.

MALDI-TOF spectrometry is excellently suited to the analysis of peptides and proteins. The analysis of nucleic acids is somewhat more difficult (Gut I G, Beck S. DNA and Matrix Assisted Laser Desorption Ionization Mass Spectrometry. *Current Innovations and Future Trends.* 1995, 1; 147-57). The sensitivity to nucleic acids is approximately 100 times worse than to peptides and decreases disproportionally with increasing fragment size. For nucleic



acids having a multiply negatively charged backbone, the ionisation process via the matrix is considerably less efficient. In MALDI-TOF spectrometry, the selection of the matrix plays an eminently important role. For the desorption of peptides, several very efficient matrixes have been found which produce a very fine crystallisation. There are now several responsive matrixes for DNA, however, the difference in sensitivity has not been reduced. The difference in sensitivity can be reduced by chemically modifying the DNA in such a manner that it becomes more similar to a peptide. Phosphorothioate nucleic acids in which the usual phosphates of the backbone are substituted with thiophosphates can be converted into a charge-neutral DNA using simple alkylation chemistry (Gut IG, Beck S. A procedure for selective DNA alkylation and detection by mass spectrometry. *Nucleic Acids Res.* 1995 Apr 25;23(8):1367-73). The coupling of a charge tag to this modified DNA results in an increase in sensitivity to the same level as that found for peptides. A further advantage of charge tagging is the increased stability of the analysis against impurities which make the detection of unmodified substrates considerably more difficult.

Genomic DNA is obtained from DNA of cell, tissue or other test samples using standard methods. This standard methodology is found in references such as Sambrook, Fritsch and Maniatis eds., *Molecular Cloning: A Laboratory Manual*, 1989.

## DESCRIPTION

Characterisation of a breast cancer in terms of its predicted aggressiveness enables the physician to make an informed decision as to a therapeutic regimen with appropriate risk and benefit trade offs to the patient.

Aggressiveness is taken to mean one or more of decreased patient survival or disease- or relapse-free survival, increased tumour-related complications and faster progression of tumour or metastases. According to the aggressiveness of the disease an appropriate treatment or treatments may be selected from the group consisting of chemotherapy, radiotherapy, surgery, biological therapy, immunotherapy, antibody treatments, treatments involving molecularly targeted drugs, estrogen receptor modulator treatments, estrogen receptor down-regulator treatments, aromatase inhibitors treatments, ovarian ablation, treatments providing LHRH analogues or other centrally acting drugs influencing estrogen production. Wherein a cancer is

characterised as 'aggressive' it is particularly preferred that a treatment such as, but not limited to, chemotherapy is provided in addition to or instead of an endocrine targeting therapy.

As used herein the term "prognostic marker" shall be taken to mean an indicator of the likelihood of progression of the disease, in particular aggressiveness and metastatic potential of a breast tumour. It is preferably used to define patients with high, low and intermediate risks of death or recurrence after treatment that result from the inherent heterogeneity of the disease process.

Indicators of tumour aggressiveness standard in the art include but are not limited to tumour stage, tumour grade, nodal status and survival.

As used herein the term "survival" shall be taken to include survival until mortality also known as overall survival (wherein said mortality may be either irrespective of cause or breast tumour related); "recurrence-free survival" (wherein the term recurrence shall include both localised and distant recurrence) ; metastasis free survival; disease free survival (wherein the term disease shall include breast cancer and diseases associated therewith). The length of said survival may be calculated by reference to a defined start point (e.g. time of diagnosis or start of treatment) and end point (e.g. death, recurrence or metastasis).

As used herein the term 'predictive marker' shall be taken to mean an indicator of response to therapy, said response is preferably defined according to patient survival. As defined herein the term predictive marker may in some situations fall within the remit of a herein described 'prognostic marker'. The two terms shall not be taken to be mutually exclusive.

Using the methods and nucleic acids described herein, statistically significant models of patient disease free survival or metastasis free survival or overall survival and/or disease progression can be developed and utilised to assist patients and clinicians in determining suitable treatment options to be included in the therapeutic regimen.

In one aspect the described method is to be used to assess the utility of therapeutic regimens comprising one or more treatments which is either an aggressive therapy such as chemotherapy or a treatment which targets the estrogen receptor pathway or is involved in estrogen metabolism, production or secretion as a therapy for patients suffering from a cell proliferative



disorder of the breast tissues. In particular this aspect of the method enables the physician to determine which treatments may be used in addition to or instead of said endocrine treatment.

In a further aspect the described method enables the characterisation of the cell proliferative disorder in terms of aggressiveness, thereby enabling the physician to recommend suitable treatments. Thus, the present invention will be seen to reduce the problems associated with present breast cell proliferative disorder treatment response prediction methods.

Using the methods and nucleic acids as described herein, patient survival can be evaluated before or during treatment for a cell proliferative disorder of the breast tissues, in order to provide critical information to the patient and clinician as to the likely progression of the disease. It will be appreciated, therefore, that the methods and nucleic acids exemplified herein can serve to improve a patient's quality of life and odds of treatment success by allowing both patient and clinician a more accurate assessment of the patient's treatment options.

The method according to the definition may be used for the improved treatment of all breast cell proliferative disorder patients, both pre and post menopausal and independent of their node or estrogen receptor status. However, it is particularly preferred that said patients are node-negative and estrogen receptor positive.

The present invention makes available a method for the improved treatment and monitoring of breast cell proliferative disorders, by enabling the accurate prediction of a patient's survival with endocrine therapy comprising one or more treatments which target the estrogen receptor pathway or are involved in estrogen metabolism, production, or secretion.

In a particularly preferred embodiment, the method according to the invention enables the differentiation between patients who have a high risk of relapse under said endocrine therapy and those who have a low risk of relapse under said therapy. The method enables the determination of a methylation pattern characteristic for a predicted survival time, in addition to the characterisation of tumours in terms of aggressiveness.

The method according to the invention may be used for the analysis of a wide variety of cell proliferative disorders of the breast tissues including, but not limited to, ductal carcinoma *in situ*, invasive ductal carcinoma, invasive lobular carcinoma, lobular carcinoma *in situ*, come-

docarcinoma, inflammatory carcinoma, mucinous carcinoma, scirrhous carcinoma, colloid carcinoma, tubular carcinoma, medullary carcinoma, metaplastic carcinoma, and papillary carcinoma and papillary carcinoma *in situ*, undifferentiated or anaplastic carcinoma and Paget's disease of the breast.

The method according to the invention is particularly suited to the prediction of survival of breast cancer in the following treatment setting. In one embodiment, the method is applied to patients who receive endocrine pathway targeting treatment as secondary treatment to an initial non chemotherapeutical therapy, e.g. surgery (hereinafter referred to as the adjuvant setting) as illustrated in Figure 1. Such a treatment is often prescribed to patients suffering from Stage 1 to 3 breast carcinomas. In this embodiment patients survival times are predicted according to their gene expression or genetic or epigenetic modifications. By detecting patients with worse disease free survival times the physician may choose to recommend the patient for further treatment, instead of or in addition to the endocrine targeting therapy(s), in particular but not limited to, chemotherapy.

The herein described invention provides a novel breast cell proliferative disorder prognostic biomarker. It is herein described that aberrant expression of the gene PITX2 is correlated to prognosis of breast cell proliferative disorder patients, in particular breast carcinoma. In particular this marker provides a novel means for the characterisation of breast carcinomas. As described herein determination of the expression of the gene PITX2 enables the prediction of survival (or outcome) of a patient treated with one or more treatments which target the estrogen receptor, synthesis or conversion pathways or are otherwise involved in estrogen metabolism, production or secretion. Survival or outcome may be based on the patient's survival or clinical or pathological tumour response, or response measured with other surrogate parameters.

The herein described invention is thereby useful for the differentiation of individuals who may be appropriately treated with one or more treatments which target the estrogen receptor pathway or are involved in estrogen metabolism, production or secretion from those individuals who would be optimally treated with other treatments in addition to said treatment. Preferred 'other treatments' include but are not limited to chemotherapy or radiotherapy.

In a further embodiment of the invention the aberrant expression of a plurality of genes comprising the gene PITX2 is analysed. Said plurality of genes is hereinafter referred to as a 'gene panel'. The analysis of multiple genes increases the accuracy of a prognosis. It is preferred that the gene panel consists of up to seven genes and/or their promoter regions associated with prognosis of breast carcinoma patients. It is further preferred that said panel consists of the gene PITX2 and one or more genes selected from the group consisting ABCA8, CDK6, ERBB2, ONECUT2, PLAUI, TBC1D3 and TFF1. It is particularly preferred that the gene panel is selected from the group of gene panels consisting of:

- PITX2, PLAUI & TFF1
- PITX2 & PLAUI
- PITX2 & TFF1

This invention therefore relates to new methods and sequences for the evaluation of adjuvant therapy of patients diagnosed with breast cell proliferative disease based on a prediction of survival or outcome.

More specifically this invention provides new methods and sequences, for patients diagnosed with breast cell proliferative disease, allowing the evaluation of adjuvant therapy, i.e. therapy before or after surgical removal of the tumour, like a cytotoxic therapy (chemotherapy) in addition to or instead of (for example in hormone receptor negative patients) an endocrine treatment, like treatment with Tamoxifen or aromatase inhibitors, wherein the evaluation is based on the prediction of the patient's survival.

One aspect of the invention is the provision of tools for predicting the survival of a patient diagnosed with a breast cell proliferative disease, such as breast cancer. These tools comprise methods for the analysis of either the expression levels of PITX2 protein, or PITX2 mRNA or the analysis of the patient's individual genetic or epigenetic modification of the gene PITX2 – summarised as the analysis of expression of the gene PITX2. Preferably the invention relates to methods for predicting the survival of a patient diagnosed with breast cancer. Preferably said patient is treated with at least one adjuvant endocrine treatment, wherein endocrine treatment is meant to comprise any treatment targeting the estrogen receptor pathway or estrogen synthesis pathway or estrogen conversion pathway i.e., which is involved in estrogen metabolism, production or secretion. Preferably the patient was treated with said adjuvant

endocrine treatment after surgical removal of the tumour. Also preferably the survival is the disease free survival.

Especially preferred are methods applied for the prediction of the disease free survival of a patient diagnosed with breast cancer under adjuvant endocrine treatment after surgical tumour removal. Even more preferred are those methods, which analyse the DNA methylation profile of the genomic region associated with the gene PITX2. Especially preferred is the analysis of the DNA methylation profile of the genomic sequence given in SEQ ID NO: 1. Especially preferred is furthermore the analysis of the methylation status of eight specific CpG dinucleotides, covered in the three sub-sequences of said SEQ ID NO: 1 given in SEQ ID NO: 13, 18 and 19. The use of nucleic acids hybridising to these nucleic acid sequences for the prediction of survival according to the invention are preferred embodiments of said invention. The use of nucleic acids hybridising to CpG positions within these nucleic acid sequences after these nucleic acids have been contacted with one or more agents that convert cytosine bases that are unmethylated at the 5'-position thereof to a base that is detectably dissimilar to cytosine in terms of hybridisation properties, for the prediction of survival according to the invention are especially preferred embodiments of said invention.

This methodology presents further improvements over the state of the art in that the method may be applied to any subject, independent of the estrogen and/or progesterone receptor status. Therefore in a preferred embodiment, the subject is not required to have been tested for estrogen or progesterone receptor status.

The object of the invention is preferably achieved by means of the analysis of the methylation pattern of PITX2 and/or its regulatory region. In a particularly preferred embodiment the sequence of said gene comprises SEQ ID NO: 1 and the sequence complementary thereto.

In one preferred embodiment the object of the invention is the prediction of survival of a subject under a treatment which targets the estrogen receptor pathway or is involved in estrogen metabolism, production or secretion. This is achieved by analysis of the expression of PITX2 and wherein it is further preferred that the sequence of said gene comprises SEQ ID NO: 1 or parts thereof.



In one aspect the invention discloses novel methods utilising the gene PITX2 for the prediction of survival of a patient diagnosed with a breast cell proliferative disease. In a preferred embodiment said patient diagnosed with a breast cell proliferative disease is treated with adjuvant endocrine monotherapy.

The invention discloses the use of the gene PITX2, as well as its promoter and regulatory elements as a prognostic marker for survival of breast cancer patients. It is preferred that these patients are treated with adjuvant endocrine monotherapy. Furthermore, the disclosed method shows the applicability of said gene to answer the question and help guiding the decision whether or not an adjuvant chemotoxic therapy shall be prescribed, preferably in addition to endocrine treatment, like the treatment with tamoxifen or aromatase inhibitors.

In one aspect of the invention, the disclosed matter provides novel nucleic acid sequences useful for the analysis of methylation within said gene, other aspects provide novel uses of the gene and the gene product as well as methods, assays and kits directed to prognosing the survival of a patient diagnosed with breast cell proliferative disease. Preferably said patient is treated with adjuvant endocrine monotherapy.

In one embodiment the method discloses the use of the gene PITX2 as a marker for the prognosis of the survival of a patient suffering from a breast cell proliferative disease. Preferably said patient is treated with adjuvant endocrine monotherapy. Said use of the gene may be enabled by means of any analysis of the expression of the gene, by means of mRNA expression analysis or protein expression analysis or by analysis of its genetic modifications leading to an altered expression (including LOH). However, in the most preferred embodiment of the invention, prediction of the survival of a patient diagnosed with breast cell proliferative disease, preferably treated with adjuvant endocrine monotherapy, is enabled by means of analysis of the methylation status of CpG sites within the gene PITX2 and its promoter or regulatory elements.

In one embodiment of the method aberrant expression of the gene PITX2 may be detected by analysis of loss of heterozygosity of the gene. In a first step genomic DNA is isolated from a biological sample of the patient's tumour. The isolated DNA is then analysed for LOH by any means standard in the art including but not limited to amplification of the gene locus or associated microsatellite markers. Said amplification may be carried out by any means standard in

the art including polymerase chain reaction (PCR), strand displacement amplification (SDA) and isothermal amplification.

The level of amplificate is then detected by any means known in the art including but not limited to gel electrophoresis and detection by probes (including Real Time PCR). Furthermore the amplificates may be labelled in order to aid said detection. Suitable detectable labels include but are not limited to fluorescence label, radioactive labels and mass labels the suitable use of which shall be described herein.

The detection of a decreased amount of an amplificate corresponding to one of the amplified alleles in a test sample as relative to that of a heterozygous control sample is indicative of LOH.

To detect the levels of mRNA encoding PITX2 in a detection system for breast cancer relapse, a sample is obtained from a patient. Said obtaining of a sample is not meant to be retrieving of a sample, as in performing a biopsy, but rather directed to the availability of an isolated biological material representing a specific tissue, relevant for the intended use. The sample can be a tumour tissue sample from the surgically removed tumour, a biopsy sample as taken by a surgeon and provided to the analyst or a sample of blood, plasma, serum or the like. The sample may be treated to extract the nucleic acids contained therein. The resulting nucleic acid from the sample is subjected to gel electrophoresis or other separation techniques. Detection involves contacting the nucleic acids and in particular the mRNA of the sample with a DNA sequence serving as a probe to form hybrid duplexes. The stringency of hybridisation is determined by a number of factors during hybridisation and during the washing procedure, including temperature, ionic strength, length of time and concentration of formamide. These factors are outlined in, for example, Sambrook et al. (Molecular Cloning: A Laboratory Manual, 2nd ed., 1989). Detection of the resulting duplex is usually accomplished by the use of labelled probes. Alternatively, the probe may be unlabeled, but may be detectable by specific binding with a ligand which is labelled, either directly or indirectly. Suitable labels and methods for labelling probes and ligands are known in the art, and include, for example, radioactive labels which may be incorporated by known methods (e.g., nick translation or kinasing), biotin, fluorescent groups, chemiluminescent groups (e.g., dioxetanes, particularly triggered dioxetanes), enzymes, antibodies, and the like.

In order to increase the sensitivity of the detection in a sample of mRNA encoding PITX2, the technique of reverse transcription/polymerisation chain reaction can be used to amplify cDNA transcribed from mRNA encoding PITX2. The method of reverse transcription/PCR is well known in the art (for example, see Watson and Fleming, *supra*).

The reverse transcription/PCR method can be performed as follows. Total cellular RNA is isolated by, for example, the standard guanidium isothiocyanate method and the total RNA is reverse transcribed. The reverse transcription method involves synthesis of DNA on a template of RNA using a reverse transcriptase enzyme and a 3' end primer. Typically, the primer contains an oligo(dT) sequence. The cDNA thus produced is then amplified using the PCR method and PITX2 specific primers. (Belyavsky et al, Nucl Acid Res 17:2919-2932, 1989; Krug and Berger, Methods in Enzymology, Academic Press, N.Y., Vol.152, pp. 316-325, 1987 which are incorporated by reference)

The present invention may also be described in certain embodiments as a kit for use in predicting the survival of a breast cancer patient before or after surgical tumour removal with or without adjuvant endocrine monotherapy state through testing of a biological sample. A representative kit may comprise one or more nucleic acid segments as described above that selectively hybridise to PITX2 mRNA and a container for each of the one or more nucleic acid segments. In certain embodiments the nucleic acid segments may be combined in a single tube. In further embodiments, the nucleic acid segments may also include a pair of primers for amplifying the target mRNA. Such kits may also include any buffers, solutions, solvents, enzymes, nucleotides, or other components for hybridisation, amplification or detection reactions. Preferred kit components include reagents for reverse transcription-PCR, in situ hybridisation, Northern analysis and/or RPA.

The present invention further provides for methods to detect the presence of the polypeptide, PITX2, in a sample obtained from a patient. It is preferred that said sequence is essentially the same as the sequence presented in SEQ ID NO: 20, as given in figure 10. Any method known in the art for detecting proteins can be used. Such methods include, but are not limited to immunodiffusion, immunoelectrophoresis, immunochemical methods, binder-ligand assays, immunohistochemical techniques, agglutination and complement assays. (for example see Basic and Clinical Immunology, Sites and Terr, eds., Appleton & Lange, Norwalk, Conn. pp 217-262, 1991 which is incorporated by reference). Preferred are binder-ligand immunoassay



methods including reacting antibodies with an epitope or epitopes of PITX2 and competitively displacing a labelled PITX2 protein or derivative thereof.

Certain embodiments of the present invention comprise the use of antibodies specific to the polypeptide encoded by the PITX2 gene. Such antibodies may be useful for prognosing the survival of a breast cancer patient preferably under adjuvant endocrine monotherapy by comparing a patient's levels of PITX2 marker expression to expression of the same marker in normal individuals. In certain embodiments production of monoclonal or polyclonal antibodies can be induced by the use of the PITX2 polypeptide as antigen. Such antibodies may in turn be used to detect expressed proteins as markers for prognosis of relapse of a breast cancer patient under adjuvant endocrine monotherapy. The levels of such proteins present in the peripheral blood of a patient may be quantified by conventional methods. Antibody-protein binding may be detected and quantified by a variety of means known in the art, such as labelling with fluorescent or radioactive ligands. The invention further comprises kits for performing the above-mentioned procedures, wherein such kits contain antibodies specific for the PITX2 polypeptides.

Numerous competitive and non-competitive protein binding immunoassays are well known in the art. Antibodies employed in such assays may be unlabeled, for example as used in agglutination tests, or labelled for use a wide variety of assay methods. Labels that can be used include radionuclides, enzymes, fluorescers, chemiluminescers, enzyme substrates or co-factors, enzyme inhibitors, particles, dyes and the like for use in radioimmunoassay (RIA), enzyme immunoassays, e.g., enzyme-linked immunosorbent assay (ELISA), fluorescent immunoassays and the like. Polyclonal or monoclonal antibodies to PITX2 or an epitope thereof can be made for use in immunoassays by any of a number of methods known in the art. One approach for preparing antibodies to a protein is the selection and preparation of an amino acid sequence of all or part of the protein, chemically synthesising the sequence and injecting it into an appropriate animal, usually a rabbit or a mouse (Milstein and Kohler *Nature* 256:495-497, 1975; Gutfre and Milstein, *Methods in Enzymology: Immunochemical Techniques* 73:1-46, Langone and Banatis eds., Academic Press, 1981 which are incorporated by reference). Methods for preparation of PITX2 or an epitope thereof include, but are not limited to chemical synthesis, recombinant DNA techniques or isolation from biological samples.



The invention provides significant improvements over the state of the art in that -at the time of filing- there are no single markers known to the public which can be used to predict the likelihood of relapse or of survival of a breast cancer patient under adjuvant endocrine monotherapy, neither from tissue samples nor from body fluid samples.

Also, no methylation marker is known which can be used to detect the likelihood of relapse or of survival of a breast cancer patient. Especially, no methylation marker is known which can be used to detect the likelihood of relapse or of survival of a breast cancer patient under adjuvant endocrine monotherapy, neither from tissue samples nor from body fluid samples.

The objective of the invention is also preferably achieved by analysis of the methylation state of the CpG dinucleotides within the genomic sequence according to SEQ ID NO: 1 and sequences complementary thereto. SEQ ID NO: 1 discloses the gene PITX2 and its promoter and regulatory elements, wherein said fragment comprises CpG dinucleotides exhibiting a disease specific methylation pattern. The methylation pattern of the gene PITX2 and its promoter and regulatory elements have heretofore not been analysed with regard to prognosis or prediction of survival of a patient diagnosed with a breast cell proliferative disorder. Due to the degeneracy of the genetic code, the sequence as identified in SEQ ID NO: 1 should be interpreted so as to include all substantially similar and equivalent sequences upstream of the promoter region of a gene which encodes a polypeptide with the biological activity of that encoded by PITX2.

In a preferred embodiment of the method, the objective of the invention is achieved by analysis of a nucleic acid comprising a sequence of at least 18 bases in length according to one of SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto.

The sequences of SEQ ID NO: 2 to 5 provide modified versions of the nucleic acid according to SEQ ID NO: 1, wherein the conversion of said sequence results in the synthesis of a nucleic acid having a sequence that is unique and distinct from SEQ ID NO: 1 as follows. (see also the following TABLE 1): SEQ ID NO: 1, sense DNA strand of PITX2 gene and its promoter and regulatory elements; SEQ ID NO: 2, converted SEQ ID NO: 1, wherein "C" converted to "T," but "CpG" remains "CpG." (i.e., corresponds to case where, for SEQ ID NO: 1, all "C" residues of CpG dinucleotide sequences are methylated and are thus not converted); SEQ ID NO: 3, complement of SEQ ID NO: 1, wherein "C" converted to "T," but "CpG"

remains "CpG" (*i.e.*, corresponds to case where, for the complement (*antisense* strand) of SEQ ID NO: 1, all "C" residues of CpG dinucleotide sequences are methylated and are thus not converted); SEQ ID NO: 4, converted SEQ ID NO: 1, wherein "C" converted to "T" for all "C" residues, including those of "CpG" dinucleotide sequences (*i.e.*, corresponds to case where, for SEQ ID NO: 1, all "C" residues of CG dinucleotide sequences are unmethylated); SEQ ID NO: 5, complement of SEQ ID NO: 1, wherein "C" converted to "T" for all "C" residues, including those of "CpG" dinucleotide sequences (*i.e.*, corresponds to case where, for the complement (*antisense* strand) of SEQ ID NO: 1, all "C" residues of CpG dinucleotide sequences are unmethylated).

**TABLE 1.** Description of SEQ ID NO: 1 to 5

SEQ ID NO: NO	Relationship to SEQ ID NO:1	Nature of cytosine base conversion
SEQ ID NO:1	Sense strand (PITX2 gene including promoter and regulatory elements)	None; untreated sequence
SEQ ID NO:2	Converted sense strand	"C" to "T," but "CpG" remains "CpG" (all "C" residues of CpGs are methylated)
SEQ ID NO:3	Converted antisense strand	"C" to "T," but "CpG" remains "CpG" (all "C" residues of CpGs are methylated)
SEQ ID NO:4	Converted sense strand	"C" to "T" for all "C" residues (all "C" residues of CpGs are <u>un</u> methylated)
SEQ ID NO:5	Converted antisense strand	"C" to "T" for all "C" residues (all "C" residues of CpGs are <u>un</u> methylated)

Significantly, heretofore, the nucleic acid sequences and molecules according to SEQ ID NO: 1 to SEQ ID NO: 5 were not implicated in or connected with the ascertainment of the prognosis of breast cancer relapse or the prediction of survival of breast cancer patients.

The described invention further discloses oligonucleotides or oligomers for detecting the cytosine methylation state within pretreated DNA, according to SEQ ID NO: 2 to SEQ ID NO: 5. The use of said oligonucleotides or oligomers comprising a nucleic acid sequence having a length of at least nine (9) nucleotides which hybridise, under moderately stringent or stringent

conditions (as defined herein above), to a pretreated nucleic acid sequence according to SEQ ID NO: 2 to SEQ ID NO: 5 and/or sequences complementary thereto is another embodiment of this invention.

Thus, the present invention includes the use of nucleic acid molecules (*e.g.*, oligonucleotides and peptide nucleic acid (PNA) molecules (PNA-oligomers)) that hybridise under moderately stringent and/or stringent hybridisation conditions to all or a portion of the sequences of SEQ ID NO: 2 to 5, or to the complements thereof for prediction of survival according to the invention. The hybridising portion of the hybridising nucleic acids is typically at least 9, 15, 20, 25, 30 or 35 nucleotides in length. However, longer molecules have inventive utility, and are thus within the scope of the present invention.

Preferably, the hybridising portion of the inventive hybridising nucleic acids is at least 95%, or at least 98%, or 100% identical to the sequence, or to a portion thereof of SEQ ID NO: 2 to 5, or to the complements thereof.

Hybridising nucleic acids of the type described herein can be used, for example, as a primer (*e.g.*, a PCR primer), or a diagnostic and/or prognostic probe or primer. Preferably, hybridisation of the oligonucleotide probe to a nucleic acid sample is performed under stringent conditions and the probe is 100% identical to the target sequence. Nucleic acid duplex or hybrid stability is expressed as the melting temperature or  $T_m$ , which is the temperature at which a probe dissociates from a target DNA. This melting temperature is used to define the required stringency conditions.

For target sequences that are related and substantially identical to the corresponding sequence of SEQ ID NO: 1 (such as PITX2 allelic variants and SNPs), rather than identical, it is useful to first establish the lowest temperature at which only homologous hybridisation occurs with a particular concentration of salt (*e.g.*, SSC or SSPE). Then, assuming that 1% mismatching results in a 1°C decrease in the  $T_m$ , the temperature of the final wash in the hybridisation reaction is reduced accordingly (for example, if sequences having > 95% identity with the probe are sought, the final wash temperature is decreased by 5°C). In practice, the change in  $T_m$  can be between 0.5°C and 1.5°C per 1% mismatch.

---

Examples of inventive oligonucleotides of length X (in nucleotides), as indicated by polynu-

cleotide positions with reference to, *e.g.*, SEQ ID NO: 1, include those corresponding to sets of consecutively overlapping oligonucleotides of length  $X$ , where the oligonucleotides within each consecutively overlapping set (corresponding to a given  $X$  value) are defined as the finite set of  $Z$  oligonucleotides from nucleotide positions:

$n$  to  $(n + (X-1))$ ;

where  $n=1, 2, 3, \dots (Y-(X-1))$ ;

where  $Y$  equals the length (nucleotides or base pairs) of SEQ ID NO: 1 ;

where  $X$  equals the common length (in nucleotides) of each oligonucleotide in the set (*e.g.*,  $X=20$  for a set of consecutively overlapping 20-mers); and

where the number ( $Z$ ) of consecutively overlapping oligomers of length  $X$  for a given SEQ ID NO: NO of length  $Y$  is equal to  $Y-(X-1)$ . For example  $Z=2,785-19=2,766$  for either sense or antisense sets of SEQ ID NO: 1, where  $X=20$ .

Preferably, the set is limited to those oligomers that comprise at least one CpG, Cpa or tpG dinucleotide, wherein 'Cpa' is indicating that said Cpa hybridises to a position (tpG) which was a CpG prior to bisulfite conversion and is a TpG now; and wherein 'tpG' is indicating that said tpG hybridises to a position (Cpa) which is the complementary to a position (tpG) which was a CpG prior to bisulfite conversion and is a TpG now.

The present invention encompasses, for *each* of SEQ ID NO: 2 to 5 (sense and antisense), the use of multiple consecutively overlapping sets of oligonucleotides or modified oligonucleotides of length  $X$ , where, *e.g.*,  $X=9, 10, 17, 20, 22, 23, 25, 27, 30$  or 35 nucleotides.

The oligonucleotides or oligomers according to the present invention constitute effective tools useful to ascertain genetic and epigenetic parameters of the genomic sequence corresponding to SEQ ID NO: 1. Preferred sets of such oligonucleotides or modified oligonucleotides of length  $X$  are those consecutively overlapping sets of oligomers corresponding to SEQ ID NO:1-5 (and to the complements thereof). Preferably, said oligomers comprise at least one CpG, tpG or Cpa dinucleotide.

Particularly preferred oligonucleotides or oligomers used to the present invention are those in which the cytosine of the CpG dinucleotide (or of the corresponding converted TpG or CpA dinucleotide) sequences is within the middle third of the oligonucleotide; that is, where the



oligonucleotide is, for example, 13 bases in length, the CpG, TpG or CpA dinucleotide is positioned within the fifth to ninth nucleotide from the 5'-end.

The oligonucleotides used in this invention can also be modified by chemically linking the oligonucleotide to one or more moieties or conjugates to enhance the activity, stability or detection of the oligonucleotide. Such moieties or conjugates include chromophores, fluorophors, lipids such as cholesterol, cholic acid, thioether, aliphatic chains, phospholipids, polyamines, polyethylene glycol (PEG), palmityl moieties, and others as disclosed in, for example, United States Patent Numbers 5,514,758, 5,565,552, 5,567,810, 5,574,142, 5,585,481, 5,587,371, 5,597,696 and 5,958,773. The probes may also exist in the form of a PNA (peptide nucleic acid) which has particularly preferred pairing properties. Thus, the oligonucleotide may include other appended groups such as peptides, and may include hybridisation-triggered cleavage agents (Krol et al., *BioTechniques* 6:958-976, 1988) or intercalating agents (Zon, *Pharm. Res.* 5:539-549, 1988). To this end, the oligonucleotide may be conjugated to another molecule, e.g., a chromophore, fluorophor, peptide, hybridisation-triggered cross-linking agent, transport agent, hybridisation-triggered cleavage agent, etc.

The oligonucleotide may also comprise at least one art-recognised modified sugar and/or base moiety, or may comprise a modified backbone or non-natural internucleoside linkage.

The oligomers used in the present invention are normally used in so called "sets" which contain at least one oligomer for analysis of each of the CpG dinucleotides of a genomic sequence comprising SEQ ID NO: 1 and sequences complementary thereto or to their corresponding CG, tG or Ca dinucleotide within the pretreated nucleic acids according to SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, wherein a 't' indicates a nucleotide which converted from a cytosine into a thymine and wherein 'a' indicates the complementary nucleotide to such a converted thymine. Preferred is a set which contains at least one oligomer for each of the CpG dinucleotides within the gene PITX2 and it's promoter and regulatory elements in both the pretreated and genomic versions of said gene, SEQ ID NO: 2 to 5 and SEQ ID NO: 1, respectively. However, it is anticipated that for economic or other factors it may be preferable to analyse a limited selection of the CpG dinucleotides within said sequences and the contents of the set of oligonucleotides should be altered accordingly. Therefore, the present invention moreover relates to a set of at least 3 n (oligonucleotides and/or PNA-oligomers) used for detecting the cytosine methylation state in pretreated geno-

mic DNA (SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto) and genomic DNA (SEQ ID NO: 1 and sequences complementary thereto). These probes enable diagnosis and/or therapy of genetic and epigenetic parameters of cell proliferative disorders. The set of oligomers may also be used for detecting single nucleotide polymorphisms (SNPs) in pretreated genomic DNA (SEQ ID NO: 2 to SEQ ID NO: 5, and sequences complementary thereto) and genomic DNA (SEQ ID NO: 1, and sequences complementary thereto) .

Moreover, the present invention includes the use of a set of at least two oligonucleotides which can be used as so-called "primer oligonucleotides" for amplifying DNA sequences of one of SEQ ID NO: 1 to SEQ ID NO: 5 and sequences complementary thereto, or segments thereof.

In the case of the sets of oligonucleotides according to the present invention, it is preferred that at least one and more preferably all members of the set of oligonucleotides is bound to a solid phase.

According to the present invention, it is preferred that an arrangement of different oligonucleotides and/or PNA-oligomers (a so-called "array") made available by the present invention is present in a manner that it is likewise bound to a solid phase. This array of different oligonucleotide- and/or PNA-oligomer sequences can be characterised in that it is arranged on the solid phase in the form of a rectangular or hexagonal lattice. The solid phase surface is preferably composed of silicon, glass, polystyrene, aluminium, steel, iron, copper, nickel, silver, or gold. However, nitrocellulose as well as plastics such as nylon which can exist in the form of pellets or also as resin matrices may also be used.

Therefore, a further subject matter of the present invention is a method for manufacturing an array fixed to a carrier material for analysis in connection with cell proliferative disorders, in which method at least one oligomer according to the present invention is coupled to a solid phase. Methods for manufacturing such arrays are known, for example, from US Patent 5,744,305 by means of solid-phase chemistry and photolabile protecting groups.

A further subject matter of the present invention relates to a DNA chip for the analysis of cell proliferative disorders. DNA chips are known, for example, in US Patent 5,837,832.

The described invention further provides a composition of matter useful for prognosing the relapse of breast cancer patients or predicting the survival of breast cancer patients. Said composition comprising at least one nucleic acid 18 base pairs in length of a segment of the nucleic acid sequence disclosed in SEQ ID NO: 2 to 5, and one or more substances taken from the group comprising :

1-5 mM Magnesium Chloride, 100-500  $\mu$ M dNTP, 0.5-5 units/10 $\mu$ l of taq polymerase, bovine serum albumen, an oligomer in particular an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to a pretreated genomic DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto. It is preferred that said composition of matter comprises a buffer solution appropriate for the stabilisation of said nucleic acid in an aqueous solution and enabling polymerase based reactions within said solution.. Suitable buffers are known in the art and commercially available.

The present invention further provides a method for conducting an assay in order to ascertain genetic and/or epigenetic parameters of the gene PITX2 and its promoter and regulatory elements. Most preferably the assay according to the following method is used in order to detect methylation within the gene PITX2 wherein said methylated nucleic acids are present in a solution further comprising an excess of background DNA, wherein the background DNA is present in between 100 to 1000 times the concentration of the DNA to be detected. Said method comprising contacting a nucleic acid sample obtained from said subject with at least one reagent or a series of reagents, wherein said reagent or series of reagents, distinguishes between methylated and non-methylated CpG dinucleotides within the target nucleic acid.

Preferably, said method comprises the following steps: In the first *step*, a sample of the tissue to be analysed is obtained. The source may be any suitable source, preferably, the source of the sample is selected from the group consisting of histological slides, biopsies, paraffin-embedded tissue, bodily fluids, plasma, serum, stool, urine, blood, nipple aspirate and combinations thereof. Preferably, the source is tumour tissue, biopsies, serum, urine, blood or nipple aspirate. The most preferred source, is the tumour sample, surgically removed from the patient or a biopsy sample of said patient.

The DNA is then isolated from the sample. Extraction may be by means that are standard to



one skilled in the art, including the use of detergent lysates, sonification and vortexing with glass beads. Once the nucleic acids have been extracted, the genomic double stranded DNA is used in the analysis.

In the second step of the method, the genomic DNA sample is treated in such a manner that cytosine bases which are unmethylated at the 5'-position are converted to uracil, thymine, or another base which is dissimilar to cytosine in terms of hybridisation behaviour. This will be understood as 'pretreatment' herein.

The above described pretreatment of genomic DNA is preferably carried out with bisulfite (hydrogen sulfite, disulfite) and subsequent alkaline hydrolysis which results in a conversion of non-methylated cytosine nucleobases to uracil or to another base which is dissimilar to cytosine in terms of base pairing behaviour. Enclosing the DNA to be analysed in an agarose matrix, thereby preventing the diffusion and renaturation of the DNA (bisulfite only reacts with single-stranded DNA), and replacing all precipitation and purification steps with fast dialysis (Olek A, et al., A modified and improved method for bisulfite based cytosine methylation analysis, *Nucleic Acids Res.* 24:5064-6, 1996) is one preferred example how to perform said pretreatment. It is further preferred that the bisulfite treatment is carried out in the presence of a radical scavenger or DNA denaturing agent.

In the third step of the method, fragments of the pretreated DNA are amplified. Wherein the source of the DNA is free DNA from serum, or DNA extracted from paraffin it is particularly preferred that the size of the amplificate fragment is between 100 and 200 base pairs in length, and wherein said DNA source is extracted from cellular sources (e.g. tissues, biopsies, cell lines) it is preferred that the amplificate is between 100 and 350 base pairs in length. It is particularly preferred that said amplificates comprise at least one 20 base pair sequence comprising at least three CpG dinucleotides. Said amplification is carried out using sets of primer oligonucleotides according to the present invention, and a preferably heat-stable polymerase. The amplification of several DNA segments can be carried out simultaneously in one and the same reaction vessel, in one embodiment of the method preferably six or more fragments are amplified simultaneously. Typically, the amplification is carried out using a polymerase chain reaction (PCR). The set of primer oligonucleotides includes at least two oligonucleotides whose sequences are each reverse complementary, identical, or hybridise under stringent or highly stringent conditions to an at least 18-base-pair long segment of the base sequences of



SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto.

In one especially preferred embodiment of the method the primers may be selected from the group consisting of SEQ ID NO: 6 to SEQ ID NO: 10.

In an alternate embodiment of the method, the methylation status of preselected CpG positions within the nucleic acid sequences comprising SEQ ID NO: 2 to SEQ ID NO: 5 may be detected by use of methylation-specific primer oligonucleotides. This technique (MSP) has been described in United States Patent No. 6,265,171 to Herman. The use of methylation status specific primers for the amplification of bisulfite treated DNA allows the differentiation between methylated and unmethylated nucleic acids. MSP primer pairs contain at least one primer which hybridises to a bisulfite treated CpG dinucleotide. Therefore, the sequence of said primers comprises at least one CpG, TpG or CpA dinucleotide. MSP primers specific for non-methylated DNA contain a "T" at the 3' position of the C position in the CpG. Preferably, therefore, the base sequence of said primers is required to comprise a sequence having a length of at least 18 nucleotides which hybridises to a pretreated nucleic acid sequence according to SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, wherein the base sequence of said oligomers comprises at least one CpG, tpG or Cpa dinucleotide. In this embodiment of the method according to the invention it is particularly preferred that the MSP primers comprise between 2 and 4 CpG, tpG or Cpa dinucleotides. It is further preferred that said dinucleotides are located within the 3' half of the primer e.g. wherein a primer is 18 bases in length the specified dinucleotides are located within the first 9 bases from the 3' end of the molecule. In addition to the CpG, tpG or Cpa dinucleotides it is further preferred that said primers should further comprise several bisulfite converted bases (i.e. cytosine converted to thymine, or on the hybridising strand, guanine converted to adenosine). In a further preferred embodiment said primers are designed so as to comprise no more than 2 cytosine or guanine bases.

The fragments obtained by means of the amplification can carry a directly or indirectly detectable label. Preferred are labels in the form of fluorescence labels, radionuclides, or detachable molecule fragments having a typical mass which can be detected in a mass spectrometer. Where said labels are mass labels, it is preferred that the labelled amplicates have a single positive or negative net charge, allowing for better detectability in the mass spectrometer. The detection may be carried out and visualised by means of, e.g., matrix assisted laser desorp-

tion/ionisation mass spectrometry (MALDI) or using electron spray mass spectrometry (ESI).

Matrix Assisted Laser Desorption/Ionization Mass Spectrometry (MALDI-TOF) is a very efficient development for the analysis of biomolecules (Karas & Hillenkamp, *Anal Chem.*, 60:2299-301, 1988). An analyte is embedded in a light-absorbing matrix. The matrix is evaporated by a short laser pulse thus transporting the analyte molecule into the vapour phase in an unfragmented manner. The analyte is ionised by collisions with matrix molecules. An applied voltage accelerates the ions into a field-free flight tube. Due to their different masses, the ions are accelerated at different rates. Smaller ions reach the detector sooner than bigger ones. MALDI-TOF spectrometry is well suited to the analysis of peptides and proteins. The analysis of nucleic acids is somewhat more difficult (Gut & Beck, *Current Innovations and Future Trends*, 1:147-57, 1995). The sensitivity with respect to nucleic acid analysis is approximately 100-times less than for peptides, and decreases disproportionally with increasing fragment size. Moreover, for nucleic acids having a multiply negatively charged backbone, the ionisation process via the matrix is considerably less efficient. In MALDI-TOF spectrometry, the selection of the matrix plays an eminently important role. For the desorption of peptides, several very efficient matrixes have been found which produce a very fine crystallisation. There are now several responsive matrixes for DNA, however, the difference in sensitivity between peptides and nucleic acids has not been reduced. This difference in sensitivity can be reduced, however, by chemically modifying the DNA in such a manner that it becomes more similar to a peptide. For example, phosphorothioate nucleic acids, in which the usual phosphates of the backbone are substituted with thiophosphates, can be converted into a charge-neutral DNA using simple alkylation chemistry (Gut & Beck, *Nucleic Acids Res.* 23: 1367-73, 1995). The coupling of a charge tag to this modified DNA results in an increase in MALDI-TOF sensitivity to the same level as that found for peptides. A further advantage of charge tagging is the increased stability of the analysis against impurities, which makes the detection of unmodified substrates considerably more difficult.

In a particularly preferred embodiment of the method the amplification of step three is carried out in the presence of at least one species of blocker oligonucleotides. The use of such blocker oligonucleotides has been described by Yu et al., *BioTechniques* 23:714-720, 1997. The use of blocking oligonucleotides enables the improved specificity of the amplification of a sub-population of nucleic acids. Blocking probes hybridised to a nucleic acid suppress, or hinder the polymerase mediated amplification of said nucleic acid. In one embodiment of the method

blocking oligonucleotides are designed so as to hybridise to background DNA. In a further embodiment of the method said oligonucleotides are designed so as to hinder or suppress the amplification of unmethylated nucleic acids as opposed to methylated nucleic acids or vice versa.

Blocking probe oligonucleotides are hybridised to the bisulfite treated nucleic acid concurrently with the PCR primers. PCR amplification of the nucleic acid is terminated at the 5' position of the blocking probe, such that amplification of a nucleic acid is suppressed where the complementary sequence to the blocking probe is present. The probes may be designed to hybridise to the bisulfite treated nucleic acid in a methylation status specific manner. For example, for detection of methylated nucleic acids within a population of unmethylated nucleic acids, suppression of the amplification of nucleic acids which are unmethylated at the position in question would be carried out by the use of blocking probes comprising a 'TpG' at the position in question, as opposed to a 'CpG.' In one embodiment of the method the sequence of said blocking oligonucleotides should be identical or complementary to molecule is complementary or identical to a sequence at least 18 base pairs in length selected from the group consisting of SEQ ID NO: 2 to 5, preferably comprising one or more CpG, TpG or CpA dinucleotides. In one embodiment of the method the sequence of said oligonucleotides is selected from the group consisting SEQ ID NO: 15 and SEQ ID NO: 16 and sequences complementary thereto.

For PCR methods using blocker oligonucleotides, efficient disruption of polymerase-mediated amplification requires that blocker oligonucleotides not be elongated by the polymerase. Preferably, this is achieved through the use of blockers that are 3'-deoxyoligonucleotides, or oligonucleotides derivatised at the 3' position with other than a "free" hydroxyl group. For example, 3'-O-acetyl oligonucleotides are representative of a preferred class of blocker molecule.

Additionally, polymerase-mediated decomposition of the blocker oligonucleotides should be precluded. Preferably, such preclusion comprises either use of a polymerase lacking 5'-3' exonuclease activity, or use of modified blocker oligonucleotides having, for example, thioate bridges at the 5'-termini thereof that render the blocker molecule nuclease-resistant. Particular applications may not require such 5' modifications of the blocker. For example, if the blocker- and primer-binding sites overlap, thereby precluding binding of the primer (e.g., with



excess blocker), degradation of the blocker oligonucleotide will be substantially precluded. This is because the polymerase will not extend the primer toward, and through (in the 5'-3' direction) the blocker - a process that normally results in degradation of the hybridised blocker oligonucleotide.

A particularly preferred blocker/PCR embodiment, for purposes of the present invention and as implemented herein, comprises the use of peptide nucleic acid (PNA) oligomers as blocking oligonucleotides. Such PNA blocker oligomers are ideally suited, because they are neither decomposed nor extended by the polymerase.

In one embodiment of the method, the binding site of the blocking oligonucleotide is identical to, or overlaps with that of the primer and thereby hinders the hybridisation of the primer to its binding site. In a further preferred embodiment of the method, two or more such blocking oligonucleotides are used. In a particularly preferred embodiment, the hybridisation of one of the blocking oligonucleotides hinders the hybridisation of a forward primer, and the hybridisation of another of the probe (blocker) oligonucleotides hinders the hybridisation of a reverse primer that binds to the amplificate product of said forward primer.

In an alternative embodiment of the method, the blocking oligonucleotide hybridises to a location between the reverse and forward primer positions of the treated background DNA, thereby hindering the elongation of the primer oligonucleotides.

It is particularly preferred that the blocking oligonucleotides are present in at least 5 times the concentration of the primers.

In the fourth step of the method, the amplificates obtained during the third step of the method are analysed in order to ascertain the methylation status of the CpG dinucleotides prior to the treatment.

In embodiments where the amplificates were obtained by means of MSP amplification and/or blocking oligonucleotides, the presence or absence of an amplificate is in itself indicative of the methylation state of the CpG positions covered by the primers and or blocking oligonucleotide, according to the base sequences thereof.. All possible known molecular biological methods may be used for this detection, including, but not limited to gel electrophoresis, se-



quencing, liquid chromatography, hybridisations, real time PCR analysis or combinations thereof. This step of the method further acts as a qualitative control of the preceding steps.

In the fourth step of the method amplicates obtained by means of both standard and methylation specific PCR are further analysed in order to determine the CpG methylation status of the genomic DNA isolated in the first step of the method. This may be carried out by means of hybridisation-based methods such as, but not limited to, array technology and probe based technologies as well as by means of techniques such as sequencing and template directed extension.

In one embodiment of the method, the amplicates synthesised in step three are subsequently hybridised to an array or a set of oligonucleotides and/or PNA probes. In this context, the hybridisation takes place in the following manner: the set of probes used during the hybridisation is preferably composed of at least 2 oligonucleotides or PNA-oligomers; in the process, the amplicates serve as probes which hybridise to oligonucleotides previously bonded to a solid phase; the non-hybridised fragments are subsequently removed; said oligonucleotides contain at least one base sequence having a length of at least 9 nucleotides which is reverse complementary or identical to a segment of the base sequences specified in the SEQ ID NO: 2 to SEQ ID NO: 5; and the segment comprises at least one CpG, TpG or CpA dinucleotide.

In a preferred embodiment, said dinucleotide is present in the central third of the oligomer. For example, wherein the oligomer comprises one CpG dinucleotide, said dinucleotide is preferably the fifth to ninth nucleotide from the 5'-end of a 13-mer. One oligonucleotide exists for the analysis of each CpG dinucleotide within the sequence according to SEQ ID NO: 1, and the equivalent positions within SEQ ID NO: 2 to 5. Said oligonucleotides may also be present in the form of peptide nucleic acids. The non-hybridised amplicates are then removed. The hybridised amplicates are detected. In this context, it is preferred that labels attached to the amplicates are identifiable at each position of the solid phase at which an oligonucleotide sequence is located.

In yet a further embodiment of the method, the genomic methylation status of the CpG positions may be ascertained by means of oligonucleotide probes that are hybridised to the bisulfite treated DNA concurrently with the PCR amplification primers (wherein said primers may either be methylation specific or standard).

A particularly preferred embodiment of this method is the use of fluorescence-based Real Time Quantitative PCR (Heid et al., *Genome Res.* 6:986-994, 1996; *also see* United States Patent No. 6,331,393). There are two preferred embodiments of utilising this method. One embodiment, known as the TaqMan™ assay employs a dual-labelled fluorescent oligonucleotide probe. The TaqMan™ PCR reaction employs the use of a non-extendible interrogating oligonucleotide, called a TaqMan™ probe, which is designed to hybridise to a CpG-rich sequence located between the forward and reverse amplification primers. The TaqMan™ probe further comprises a fluorescent "reporter moiety" and a "quencher moiety" covalently bound to linker moieties (*e.g.*, phosphoramidites) attached to the nucleotides of the TaqMan™ oligonucleotide. Hybridised probes are displaced and broken down by the polymerase of the amplification reaction thereby leading to an increase in fluorescence. For analysis of methylation within nucleic acids subsequent to bisulfite treatment, it is required that the probe be methylation specific, as described in United States Patent No. 6,331,393, (hereby incorporated by reference in its entirety) also known as the MethyLight assay. The second preferred embodiment of this MethyLight technology is the use of dual-probe technology (Lightcycler®), each probe carrying donor or recipient fluorescent moieties, hybridisation of two probes in proximity to each other is indicated by an increase or fluorescent amplification primers. Both these techniques may be adapted in a manner suitable for use with bisulfite treated DNA, and moreover for methylation analysis within CpG dinucleotides.

Also any combination of these probes or combinations of these probes with other known probes may be used.

In a further preferred embodiment of the method, the fourth step of the method comprises the use of template-directed oligonucleotide extension, such as MS-SNuPE as described by Gonzalgo & Jones, *Nucleic Acids Res.* 25:2529-2531, 1997. In said embodiment it is preferred that the methylation specific single nucleotide extension primer (MS-SNuPE primer) is identical or complementary to a sequence at least nine but preferably no more than twenty five nucleotides in length of one or more of the sequences taken from the group of SEQ ID NO: 2 to SEQ ID NO: 5. However it is preferred to use fluorescently labelled nucleotides, instead of radiolabelled nucleotides.

In yet a further embodiment of the method, the fourth step of the method comprises sequenc-

ing and subsequent sequence analysis of the amplificate generated in the third step of the method (Sanger F., et al., *Proc Natl Acad Sci USA* 74:5463-5467, 1977).

Additional embodiments of the invention provide a method for the analysis of the methylation status of genomic DNA according to the invention (SEQ ID NO: 1) without the need for pre-treatment.

In the *first step* of such additional embodiments, the genomic DNA sample is isolated from tissue or cellular sources. Preferably, such sources include cell lines, histological slides, biopsy tissue, body fluids, or breast tumour tissue embedded in paraffin. Extraction may be by means that are standard to one skilled in the art, including but not limited to the use of detergent lysates, sonification and vortexing with glass beads. Once the nucleic acids have been extracted, the genomic double-stranded DNA is used in the analysis.

In a preferred embodiment, the DNA may be cleaved prior to the treatment, and this may be by any means standard in the state of the art, but preferably with methylation-sensitive restriction endonucleases.

In the *second step*, the DNA is then digested with one or more methylation sensitive restriction enzymes. The digestion is carried out such that hydrolysis of the DNA at the restriction site is informative of the methylation status of a specific CpG dinucleotide.

In the *third step*, which is optional but a preferred embodiment, the restriction fragments are amplified. This is preferably carried out using a polymerase chain reaction, and said amplificates may carry suitable detectable labels as discussed above, namely fluorophore labels, radionuclides and mass labels.

In the *final step* the amplificates are detected. The detection may be by any means standard in the art, for example, but not limited to, gel electrophoresis analysis, hybridisation analysis, incorporation of detectable tags within the PCR products, DNA array analysis, MALDI or ESI analysis.

The present invention enables prognosis of events which are disadvantageous to patients or individuals in which important genetic and/or epigenetic parameters within the PITX2 gene



and its promoter or regulatory elements may be used as prognostic markers for breast cancer relapse or as 'adjuvant marker' for prediction of need of additional treatment besides of endocrine monotherapy. Said parameters obtained by means of the present invention may be compared to another set of genetic and/or epigenetic parameters, the differences serving as the basis for a prognosis of events which are disadvantageous to patients or individuals.

Specifically, the present invention provides for prognostic cancer relapse assays based on measurement of differential methylation of PITX2 CpG dinucleotide sequences. Preferred gene sequences useful to measure such differential methylation are represented herein by SEQ ID NO: 1 to 5. Typically, such assays involve obtaining a tissue sample from a test tissue, performing an assay to measure the methylation status of at least one of the inventive PITX2-specific CpG dinucleotide sequences derived from the tissue sample, relative to a control sample, and making a diagnosis or prognosis or prediction based thereon.

In particular preferred embodiments, inventive oligomers to assess PITX2 specific CpG dinucleotide methylation status, such as those based on SEQ ID NO: 1 to 5, or arrays thereof, as well as a kit based thereon are used for the prognosis of breast cancer relapse and/or the prediction of survival of a patient diagnosed with breast cancer, preferably under endocrine treatment since surgical removal of the tumour.

Moreover, an additional aspect of the present invention is a kit comprising, for example: a bisulfite-containing reagent as well as at least one oligonucleotide whose sequences in each case correspond, are complementary, or hybridise under stringent or highly stringent conditions to a 18-base long segment of the sequences SEQ ID NO: 1 to 5. Said kit may further comprise instructions for carrying out and evaluating the described method. In a further preferred embodiment, said kit may further comprise standard reagents for performing a CpG position-specific methylation analysis, wherein said analysis comprises one or more of the following techniques: MS-SNuPE, MSP, MethyLight™, HeavyMethyl™, COBRA, and nucleic acid sequencing. However, a kit along the lines of the present invention can also contain only part of the aforementioned components.

Typical reagents (*e.g.*, as might be found in a typical COBRA-based kit) for COBRA analysis may include, but are not limited to: PCR primers for specific gene (or methylation-altered DNA sequence or CpG island); restriction enzyme and appropriate buffer; gene-hybridisation



oligo; control hybridisation oligo; kinase labelling kit for oligo probe; and radioactive nucleotides. Additionally, bisulfite conversion reagents may include: DNA denaturation buffer; sulfonation buffer; DNA recovery reagents or kits (*e.g.*, precipitation, ultrafiltration, affinity column); desulfonation buffer; and DNA recovery components.

Typical reagents (*e.g.*, as might be found in a typical MethyLight®-based kit) for MethyLight® analysis may include, but are not limited to: PCR primers for specific gene (or methylation-altered DNA sequence or CpG island); TaqMan® probes; optimised PCR buffers and deoxynucleotides; and Taq polymerase.

Typical reagents (*e.g.*, as might be found in a typical Ms-SNuPE-based kit) for Ms-SNuPE analysis may include, but are not limited to: PCR primers for specific gene (or methylation-altered DNA sequence or CpG island); optimised PCR buffers and deoxynucleotides; gel extraction kit; positive control primers; Ms-SNuPE primers for specific gene; reaction buffer (for the Ms-SNuPE reaction); and radioactive nucleotides. Additionally, bisulfite conversion reagents may include: DNA denaturation buffer; sulfonation buffer; DNA recovery reagents or kit (*e.g.*, precipitation, ultrafiltration, affinity column); desulfonation buffer; and DNA recovery components.

Typical reagents (*e.g.*, as might be found in a typical MSP-based kit) for MSP analysis may include, but are not limited to: methylated and unmethylated PCR primers for specific gene (or methylation-altered DNA sequence or CpG island), optimised PCR buffers and deoxynucleotides, and specific probes.

Specifically, the present invention is related to a method for characterising a cell proliferative disorder of the breast tissues and/or predicting the survival of a patient diagnosed with said disorder, comprising the steps of: a) detecting the expression of a nucleic acid or a polypeptide expressed from the PITX2 gene in an isolated biological sample representative of said cell proliferative disorders of the breast tissues and b) predicting therefrom the survival of said patient, characteristics of said cell proliferative disorder, and/or prognosis of said patient. Preferably, the method according to the present invention further comprises c) determining a suitable treatment regimen for the subject.

---

Preferred is a method according to the present invention, wherein the patient is characterised

by being subject to adjuvant endocrine therapy comprising one or more treatments which target the estrogen receptor pathway or are involved in estrogen metabolism, production or secretion.

Preferred is also a method according to the present invention, wherein said breast cell proliferative disorders are taken from the group comprising ductal carcinoma *in situ*, invasive ductal carcinoma, invasive lobular carcinoma, lobular carcinoma *in situ*, comedocarcinoma, inflammatory carcinoma, mucinous carcinoma, scirrhous carcinoma, colloid carcinoma, tubular carcinoma, medullary carcinoma, metaplastic carcinoma, and papillary carcinoma and papillary carcinoma *in situ*, undifferentiated or anaplastic carcinoma and Paget's disease of the breast.

According to another aspect of the method according to the present invention, said method is characterised in that the detection is carried out by a) amplification of a section of the gene PITX2 and/or microsatellites associated therewith; b) detecting the presence and/or absence of alleles of the amplificate; and c) predicting therefrom the survival of said patient, characteristics of said cell proliferative disorder, and/or prognosis of said patient.

According to another aspect of the method according to the present invention, said method is characterised in that the detection is carried out by a) contacting said biological sample with an antibody immunoreactive with the PITX2 polypeptide to form an immunocomplex; b) detecting said immunocomplex; and c) predicting therefrom the survival of said patient, characteristics of said cell proliferative disorder, and/or prognosis of said patient.

According to another aspect of the method according to the present invention, said method is characterised in that the detection is carried out by a) contacting said biological sample with an antibody immunoreactive with the PITX2 polypeptide to form an immunocomplex; b) detecting said immunocomplex; c) therefrom predicting the survival of said patient, characteristics of said cell proliferative disorder, and/or prognosis of said patient, and d) comparing the quantity of said immunocomplex to the quantity of immunocomplex formed under identical conditions with the same antibody and a control sample from one or more patients with a known prognosis.

According to yet another aspect of the method according to the present invention, the detection is carried out by a) contacting said biological sample with an antibody immunoreactive with the PITX2 polypeptide to form an immunocomplex; b) detecting said immunocomplex; and c) predicting therefrom the survival of said patient, characteristics of said cell proliferative disorder, and/or prognosis of said patient, and wherein an increase in quantity of said immunocomplex in the sample from said subject relative to said control sample is indicative of a bad prognosis.

Preferred is a method according to the present invention, wherein said immunoassay is a radioimmunoassay or an ELISA or a Western blot. Preferred is a method according to the present invention, wherein said detection is afforded by mRNA expression analysis. Most preferred is a method according to the present invention, comprising detecting the level of mRNA encoding a PITX2 polypeptide in a biological sample from a patient,

Preferred is a method according to the present invention, wherein a increased concentration of said mRNA above the concentration determined for an individual known to have a good prognosis indicates a bad prognosis.

According to yet another aspect of the method according to the present invention said method comprises the steps of: a) providing a polynucleotide probe which specifically hybridises or is identical to a polynucleotide consisting of SEQ ID NO: 19 or SEQ ID NO: 1, b) incubating said sample with said polynucleotide probe under high stringency conditions to form a specific hybridisation complex between a nucleic acid and said probe; and c) detecting said hybridisation complex.

Preferred is a method according to the present invention wherein said nucleic acid is mRNA or a cDNA derived therefrom.

Preferred is a method according to the present invention wherein the detecting step further comprises the steps of: a) producing a cDNA from mRNA in the sample; b) providing two oligonucleotides which specifically hybridise to regions flanking a segment of the cDNA; c) performing a polymerase chain reaction on the cDNA of step a) using the oligonucleotides of step b) as primers to amplify the cDNA segment; and d) detecting the amplified cDNA segment.

Yet another aspect of the present invention relates to a use of a polypeptide expressed from the PITX2 gene for differentiating or distinguishing between patients diagnosed with breast cancer, who have a good survival prognosis and patients who have a bad survival prognosis. Preferably, said polypeptide is expressed from the PITX2 gene and used for prediction of survival of a patient diagnosed with a cell proliferative disorder of the breast.

Preferred is a method according to the present invention wherein said detection comprises determining the genetic parameters of the gene PITX2, its promoter and/or regulatory elements. More preferred is a method according to the present invention wherein said detection comprises determining the epigenetic parameters of the gene PITX2, its promoter and/or regulatory elements.

Preferred is a method according to the present invention, wherein said detection comprises determining the methylation status of one or more CpG positions of a target nucleic acid within the gene PITX2, its promoter and/or regulatory elements, in particular through the methylation analysis of a genomic DNA sequence according to SEQ ID NO: 1. Preferred is further a method according to the present invention, wherein said detection comprises determining the methylation status of one or more CpG positions of a target nucleic acid characterised as being identical to or hybridising under stringent or moderately stringent conditions to a sequence out of the group of SEQ ID NO: NOs 13, 18 and 19. Preferred is further a method according to the present invention, wherein the methylation analysis is afforded by contacting said target nucleic acid with one or more agents that convert cytosine bases that are unmethylated at the 5'-position thereof to a base that is detectably dissimilar to cytosine in terms of hybridisation properties.

Preferred is a method according to the present invention, wherein contacting said target nucleic acids with one or more agents comprises use of a solution selected from the group consisting of bisulfite, hydrogen sulfite, disulfite, and combinations thereof.

Yet another aspect of the present invention relates to the use of a set of oligomer probes comprising at least two oligomers, in particular an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or



stringent conditions to a pretreated genomic DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for detecting the cytosine methylation state and/or single nucleotide polymorphisms (SNPs) within one of the sequences according to SEQ ID NO: 1, and sequences complementary thereto.

Yet another aspect of the present invention relates to a method for manufacturing an arrangement of different oligomers (array) fixed to a carrier material for analysing diseases associated with the methylation state of the CpG dinucleotides of one of SEQ ID NO: 1, and sequences complementary thereto wherein at least one oligomer, in particular an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to a pretreated genomic DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, is coupled to a solid phase.

Yet another aspect of the present invention relates to a composition of matter comprising the following: a) a nucleic acid comprising a sequence at least 18 bases in length of a segment of the chemically pretreated genomic DNA according to one of the sequences taken from the group comprising SEQ ID NO: 1 to SEQ ID NO: 5 and sequences complementary thereto, and b) a buffer comprising at least one of the following substances: 1 to 5 mM magnesium chloride, 100-500  $\mu$ M dNTP, 0.5-5 units/10ul of taq polymerase, an oligomer, in particular an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to a pretreated genomic DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto.

Preferably, the gene PITX2, its promoter and/or regulatory elements is used for predicting the survival of patients diagnosed with a cell proliferative disease. Preferred is the use of the mRNA of the gene PITX2 for predicting the survival of patients diagnosed with a cell proliferative disease.

Yet another aspect of the present invention relates to a method for predicting the survival of patients diagnosed with a cell proliferative disease according to the present invention, com-

prising: a) isolating or enriching genomic DNA from said biological sample ; b) treating the genomic DNA, or a fragment thereof, with one or more reagents to convert 5-position unmethylated cytosine bases to uracil or to another base that is detectably dissimilar to cytosine in terms of hybridisation properties; c) contacting the treated genomic DNA, or the treated fragment thereof, with an amplification enzyme and at least two primers comprising, in each case a contiguous sequence at least 18 nucleotides in length that is complementary to, or hybridises under moderately stringent or stringent conditions to a sequence selected from the group consisting of SEQ ID NO: 2 to 5, and complements thereof, wherein the treated DNA or a fragment thereof is either amplified to produce one or more amplicates, or is not amplified; and d) determining, based on the presence or absence of, or on the quantity or on a property of said amplicate, the methylation state of at least one CpG dinucleotide sequence of SEQ ID NO: 1, or an average, or a value reflecting an average methylation state of a plurality of CpG dinucleotide sequences of SEQ ID NO: 1.

Yet another aspect of the present invention relates to a method for detecting the survival of patients diagnosed with a cell proliferative disease of the breast according to the present invention, comprising the following steps of a) obtaining, from a subject, a biological sample having subject genomic DNA; b) treating the genomic DNA, or a fragment thereof, with one or more reagents to convert 5-position unmethylated cytosine bases to uracil or to another base that is detectably dissimilar to cytosine in terms of hybridisation properties; c) amplifying one or more fragments of the treated DNA such that only DNA originating from breast or breast cell proliferative disorder cells are amplified and d) detecting the amplicates or characteristics thereof and thereby deducing on the survival of patients diagnosed with a cell proliferative disease of the breast.

Yet another aspect of the present invention relates to a use of an oligomer, an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to an artificially modified, chemically pretreated DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for differentiating or distinguishing between patients diagnosed with breast cancer, who have a good survival prognosis and patients who have a bad survival prognosis.

Yet another aspect of the present invention relates to a use of a nucleic acid comprising a sequence of at least 18 bases in length of a segment of the artificially modified, chemically pretreated, DNA according to one of the sequences taken from the group comprising SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for differentiating or distinguishing between patients diagnosed with breast cancer, who have a good survival prognosis and patients who have a bad survival prognosis. Yet another aspect of the present invention relates to a use of an oligomer, an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to an artificially modified, chemically pretreated DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for prediction of survival of a patient diagnosed with a cell proliferative disorder of the breast.

Yet another aspect of the present invention relates to a use of a nucleic acid comprising a sequence of at least 18 bases in length of a segment of the artificially modified, chemically pretreated, DNA according to one of the sequences taken from the group comprising SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for prediction of survival of a patient diagnosed with a cell proliferative disorder of the breast. Yet another aspect of the present invention relates to a use of a nucleic acid represented by their SEQ ID NO: NO out of the group of nucleic acids according to SEQ ID NO: NOS : 6, 7, 8, 9, 10, 14, 15, 22, 23, 24, 25, 26, 27, 28, 29, 30 and 31, for differentiating or distinguishing between patients diagnosed with breast cancer, who have a good survival prognosis and patients who have a bad survival prognosis. Yet another aspect of the present invention relates to a use of a nucleic acid represented by their SEQ ID NO: NO out of the group of nucleic acids according to SEQ ID NO: NOS : 6, 7, 8, 9, 10, 14, 15, 22, 23, 24, 25, 26, 27, 28, 29, 30 and 31, for prediction of survival of a patient diagnosed with a cell proliferative disorder of the breast.

In the context of this invention the terms "obtaining a biological sample" or "obtaining a sample from a subject", is supposed to comprise several different sources of such a sample, but always excludes the active retrieval of a sample from an individual patient, such as the performance of a biopsy. Included are the following examples: obtaining a sample, which was prior to the obtaining step taken from a patient in a biopsy or surgery, from a sample provider; obtaining a sample from a clinician, a surgeon or other medical personnel; obtaining a sample from a courier, who is bringing the sample from the clinician or practitioner or patient him-



self, for example to the analytic service station, as well as obtaining a sample such as a body fluid sample per post or from the hands of the patient himself. This is not meant to be a limiting list, but shall illustrate that it is not a feature of the invention that it needs to be carried out on the patient itself.

The term "biological material" relates to any material that is derived from a source, in particular an animal and/or human source, that contains or is suspected to contain genomic DNA. On example of a biological material according to the present invention will be a biological sample.

In the context of the present invention, the term "CpG island" refers to a contiguous region of genomic DNA that satisfies the criteria of (1) having a frequency of CpG dinucleotides corresponding to an "Observed/Expected Ratio"  $>0.6$ , and (2) having a "GC Content"  $>0.5$ . CpG islands are typically, but not always, between about 0.2 to about 1 kb in length.

In the context of the present invention the term "regulatory region" of a gene is taken to mean nucleotide sequences which affect the expression of a gene. Said regulatory regions may be located within, proximal or distal to said gene. Said regulatory regions include but are not limited to constitutive promoters, tissue-specific promoters, developmental-specific promoters, inducible promoters and the like. Promoter regulatory elements may also include certain enhancer sequence elements that control transcriptional or translational efficiency of the gene.

In the context of the present invention, the term "methylation" refers to the presence or absence of 5-methylcytosine ("5-mCyt") at one or a plurality of CpG dinucleotides within a DNA sequence.

In the context of the present invention the term "methylation state" is taken to mean the degree of methylation present in a nucleic acid of interest, this may be expressed in absolute or relative terms i.e. as a percentage or other numerical value or by comparison to another tissue and therein described as hypermethylated, hypomethylated or as having significantly similar or identical methylation status.

In the context of the present invention, the term "hemi-methylation" or "hemimethylation" refers to the methylation state of a palindromic CpG methylation site, where only a single



cytosine in one of the two CpG dinucleotide sequences of the double stranded CpG methylation site is methylated (e.g., 5'-NNC<sup>M</sup>GNN-3' (top strand): 3'-NNGCNN-5' (bottom strand)).

In the context of the present invention, the term "hypermethylation" refers to the average methylation state corresponding to an *increased* presence of 5-mCyt at one or a plurality of CpG dinucleotides within a DNA sequence of a test DNA sample, relative to the amount of 5-mCyt found at corresponding CpG dinucleotides within a normal control DNA sample.

In the context of the present invention, the term "hypomethylation" refers to the average methylation state corresponding to a *decreased* presence of 5-mCyt at one or a plurality of CpG dinucleotides within a DNA sequence of a test DNA sample, relative to the amount of 5-mCyt found at corresponding CpG dinucleotides within a normal control DNA sample.

In the context of the present invention, the term "microarray" refers broadly to both "DNA microarrays," and "DNA chip(s)," as recognised in the art, encompasses all art-recognised solid supports, and encompasses all methods for affixing nucleic acid molecules thereto or synthesis of nucleic acids thereon.

"Genetic parameters" are mutations and polymorphisms of genes and sequences further required for their regulation. To be designated as mutations are, in particular, insertions, deletions, point mutations, inversions and polymorphisms and, particularly preferred, SNPs (single nucleotide polymorphisms).

"Epigenetic modifications" or "epigenetic parameters" are modifications of DNA bases of genomic DNA and sequences further required for their regulation, in particular, cytosine methylations thereof. Further epigenetic parameters include, for example, the acetylation of histones which, however, cannot be directly analysed using the described method but which, in turn, correlate with the DNA methylation.

In the context of the present invention, the term "bisulfite reagent" refers to a reagent comprising bisulfite, disulfite, hydrogen sulfite or combinations thereof, useful as disclosed herein to distinguish between methylated and unmethylated CpG dinucleotide sequences.

In the context of the present invention, the term "Methylation assay" refers to any assay for determining the methylation state of one or more CpG dinucleotide sequences within a sequence of DNA.

In the context of the present invention, the term "MS.AP-PCR" (Methylation-Sensitive Arbitrarily-Primed Polymerase Chain Reaction) refers to the art-recognised technology that allows for a global scan of the genome using CG-rich primers to focus on the regions most likely to contain CpG dinucleotides, and described by Gonzalgo et al., *Cancer Research* 57:594-599, 1997.

In the context of the present invention, the term "MethyLight®" refers to the art-recognised fluorescence-based real-time PCR technique described by Eads et al., *Cancer Res.* 59:2302-2306, 1999.

In the context of the present invention, the term "HeavyMethyl™" assay, in the embodiment thereof implemented herein, refers to a HeavyMethyl™ MethyLight assay, which is a variation of the MethyLight assay, wherein the MethyLight assay is combined with methylation specific *blocking* probes covering CpG positions between the amplification primers.

The term "Ms-SNuPE" (Methylation-sensitive Single Nucleotide Primer Extension) refers to the art-recognised assay described by Gonzalgo & Jones, *Nucleic Acids Res.* 25:2529-2531, 1997.

In the context of the present invention the term "MSP" (Methylation-specific PCR) refers to the art-recognised methylation assay described by Herman et al. *Proc. Natl. Acad. Sci. USA* 93:9821-9826, 1996, and by US Patent No. 5,786,146.

In the context of the present invention the term "COBRA" (Combined Bisulfite Restriction Analysis) refers to the art-recognised methylation assay described by Xiong & Laird, *Nucleic Acids Res.* 25:2532-2534, 1997.

In the context of the present invention the term "hybridisation" is to be understood as a bond of an oligonucleotide to a complementary sequence along the lines of the Watson-Crick base pairings in the sample DNA, forming a duplex structure.

"Stringent hybridisation conditions," as defined herein, involve hybridising at 68°C in 5x SSC/5x Denhardt's solution/1.0% SDS, and washing in 0.2x SSC/0.1% SDS at room temperature, or involve the art-recognised equivalent thereof (e.g., conditions in which a hybridisation is carried out at 60°C in 2.5 x SSC buffer, followed by several washing steps at 37°C in a low buffer concentration, and remains stable). Moderately stringent conditions, as defined herein, involve including washing in 3x SSC at 42°C, or the art-recognised equivalent thereof. The parameters of salt concentration and temperature can be varied to achieve the optimal level of identity between the probe and the target nucleic acid. Guidance regarding such conditions is available in the art, for example, by Sambrook et al., 1989, *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, N.Y.; and Ausubel et al. (eds.), 1995, *Current Protocols in Molecular Biology*, (John Wiley & Sons, N.Y.) at Unit 2.10.

"Background DNA" as used herein refers to any nucleic acids which originate from sources other than colon cells.

In the context of this application "survival" is meant to describe the time from diagnosis or start of treatment to an endpoint, which may be either the time of death (considering any reason for death or only death from breast cancer), or the time of recurrence of breast cancer (for example in form of metastases), which may be local or distant, or the time of occurrence of any breast cancer associated disease. Therefore "predicting the survival" is meant to comprise predicting the disease free survival, as well as the overall survival or any other consideration of time between diagnosis and endpoint of treatment. However, as it is obvious in the state of the art, a precise prediction of life time is generally impossible whether it is based on a biomarker analysis, or any other prognostic tools, it is understood throughout the invention that said term "prediction of survival" or "predicting the survival" is used to describe the risk of patient to suffer from a recurrence of metastasis or other disease caused by the original breast cell proliferative disease the patient was diagnosed with (also termed "risk of relapse"). Said risk can be predicted with a certain probability or likelihood. It is also clear, that predicting the survival is meant to comprise the determination of the likelihood or probability whether a subject or patient will survive for a longer or shorter period of time.

Throughout this invention it is preferred that said survival is characterised as the disease free or the overall survival. It is especially preferred that survival is understood as disease free

survival. Disease free survival is understood as absence of recurrence of cancer (local or distant).

The terms "endocrine therapy" or "endocrine treatment" is meant to comprise any therapy, treatment or treatments targeting the estrogen receptor pathway or estrogen synthesis pathway or estrogen conversion pathway, which is involved in estrogen metabolism, production or secretion. Said treatments include, but are not limited to estrogen receptor modulators, estrogen receptor down-regulators, aromatase inhibitors, ovarian ablation, LHRH analogues and other centrally acting drugs influencing estrogen production.

The term "monotherapy" is used to explain that no other treatment is given in addition or to support said monotherapy.

In the context of the present invention the term "chemotherapy" is taken to mean the use of drugs or chemical substances to treat cancer. This definition excludes radiation therapy (treatment with high energy rays or particles), hormone therapy (treatment with hormones or hormone analogues (synthetic substitutes) and surgical treatment.

In the context of the present invention the term "adjuvant treatment" is taken to mean a therapy of a cancer patient immediately following an initial non chemotherapeutical therapy, e.g. surgery. In general, the purpose of an adjuvant therapy is to provide a significantly smaller risk of recurrences compared without the adjuvant therapy.

In the context of the present invention the term "determining a suitable treatment regimen for the subject" is taken to mean a treatment regimen (i.e. a single therapy or a combination of different therapies that are used for the prevention and/or treatment of the cancer in the patient) for the cancer patient that is started, modified and/or ended based or essentially based or at least partially based on the results of the analysis according to the present invention. One example is starting an adjuvant endocrine therapy after surgery, another would be to modify the dosage of a particular chemotherapy. The determination can, in addition to the results of the analysis according to the present invention, be based on personal characteristics of the subject to be treated. In most cases, the actual determination of the suitable treatment regimen for the subject will be performed by the attending physician or doctor.



While the present invention has been described with specificity in accordance with certain of its preferred embodiments, the following examples and figures serve only to illustrate the invention and is not intended to limit the invention within the principles and scope of the broadest interpretations and equivalent configurations thereof.

In the accompanying sequence protocol and the figures,

SEQ ID NO: 1 shows the nucleic acid sequence of the human gene PITX2,

SEQ ID NO: 2 to 5 show chemically pretreated nucleic acid sequences of the gene PITX2, according to table 1.

SEQ ID NO: 6 to 9 show the nucleic acid sequences of those primers and probes useful to predict the survival of breast cancer patients according to the invention as described in example 4.

SEQ ID NO: 14 to 17 show the nucleic acid sequences of those primers and probes useful to predict the survival of breast cancer patients according to the invention as described in example 5.

SEQ ID NO: 10 to 12 show the nucleic acid sequences of primers and probes according to a control gene used in the example 4 and 5.

SEQ ID NO: 13 shows a subsequence of SEQ ID NO: 1, which represents the nucleic acid sequence of the human gene PITX2.

SEQ ID NO: 18 shows an amino acid sequence of the polypeptide encoded by the gene PITX2. The amino acid sequence of the polypeptide encoded by the gene PITX2 is also illustrated in figure 10.

## FIGURES

Figure 1 presents a scheme to illustrate a preferred application of the method according to the invention. Along the Y axis tumour(s) mass (or size) increases, wherein the line '3' indicates the limit of detectability of said tumour mass. The X axis represents time (such as in life time of a patient). Accordingly said figure illustrates a simplified model of an Stage 1-3 breast tumour wherein primary treatment was surgery (at point 1), followed by adjuvant therapy with Tamoxifen, as an example for an endocrine treatment. In a first scenario a patient without relapse during endocrine treatment (4) is shown as remaining below the limit of detectability for the duration of the observation. A patient with relapse of the cancer (5) has a period of disease free survival (2) followed by relapse when the carcinoma mass reaches the level of detectability.

Figure 2 shows the result of the assay (QM assay) as described in example 4: A Kaplan-Meier estimated metastasis-free survival curve for 3 CpG sites of the PITX2 gene by means of Real-Time methylation specific probe analysis (QM assay). The lower curve shows the proportion of metastasis free patients in the population with above median methylation levels, the upper curve shows the proportion of metastasis free patients in the population with below median methylation levels. The X axis shows the metastasis free survival times of the patients in months, and the Y axis shows the proportion of metastasis free survival patients.

Figure 3 shows the result of the chip hybridisation experiment as described in example 2. A Kaplan-Meier estimated metastasis-free survival curves for 2 CpG positions of the PITX2 gene by means of methylation specific detection oligo hybridisation analysis. The lower curve shows the proportion of metastasis free patients in the population with above median methylation levels, the upper curve shows the proportion of metastasis free patients in the population with below median methylation levels. The X axis shows the metastasis free survival times of the patients in months, and the Y axis shows the proportion of metastasis free survival patients.

Figure 4 shows the Kaplan-Meier estimated metastasis-free survival curves for 2 CpG positions of the PITX2 gene by means of methylation specific detection oligo hybridisation analysis. The lower line shows the proportion of metastasis free patients in the population of 55 patients with above median methylation levels, the upper curve shows the proportion of metastasis free patients in the population of 54 patients with below median methylation levels. The X axis shows the metastasis free survival times of the patients in years, and the Y axis shows the proportion of metastasis free survival patients in %. This resulted from a first data set that was achieved in a first study.

Figure 5 shows the Kaplan-Meier estimated metastasis-free survival curves for 6 different CpG positions located within the preferred region of the PITX2 gene (SEQ ID NO: 13) by means of methylation specific detection oligo hybridisation analysis. The lower line shows the proportion of metastasis free patients in the population of 118 patients with above median methylation levels, the upper curve shows the proportion of metastasis free patients in the population of 118 patients with below median methylation levels. The X axis shows the metastasis free survival times of the patients in years, and the Y axis shows the proportion of

metastasis free survival patients in %. This resulted from a second data set that was achieved in a second study.

Figure 6 shows the Kaplan-Meier estimated metastasis-free survival curves for 6 different CpG positions located within the preferred region of the PITX2 gene (SEQ ID NO: 13) by means of methylation specific detection oligo hybridisation analysis. This time only a subpopulation of 148 patients, characterised by a tumour at grade G1 or G2, was analysed: The lower curve shows the proportion of metastasis free patients in the population of 74 patients with above median methylation levels, the upper curve shows the proportion of metastasis free patients in the population of 74 patients with below median methylation levels. The X axis shows the metastasis free survival times of the patients in years, and the Y axis shows the proportion of metastasis free survival patients in %. This resulted from a second data set that was achieved in the second example.

Figure 7 shows the Kaplan-Meier estimated metastasis-free survival curves for 4 different CpG positions located within the preferred region of the PITX2 gene (SEQ ID NO: 13) by means of methylation specific detection oligo hybridisation analysis. This time a subpopulation of 224 patients, characterised by a tumour of stage 1 or 2 (T1 or T2), was analysed: The lower curve shows the proportion of metastasis free patients in the population of 112 patients with above median methylation levels, the upper curve shows the proportion of metastasis free patients in the population of 112 patients with below median methylation levels. The X axis shows the metastasis free survival times of the patients in years, and the Y axis shows the proportion of metastasis free survival patients in %. This resulted from the second data set that was achieved in the second example.

Figure 8 shows the disease-free survival curves for a combination of two oligonucleotides each from the genes TBC1D3 and CDK6, and one oligonucleotide from the gene PITX2 covering two CpG sites. The black curve shows the proportion of disease free patients in the population with above median methylation scores, the grey curve shows shows the proportion of disease free patients in the population with below median methylation scores.

Figure 9 shows the plot according to Figure 8 and the classification of the sample set by means of the St. Gallen method. The unbroken lines represent the methylation analysis wherein the black curve shows the proportion of disease free patients in the population with



above median methylation scores, the grey curve shows the proportion of disease free patients in the population with below median methylation scores. The broken lines represent the St. Gallen classification of the sample set wherein the black curve shows the disease free survival time of the high risk group and the grey curve shows the disease free survival of the low risk group.

Figure 10 illustrates the amino acid sequence of the polypeptide encoded by the gene PITX2.

## EXAMPLES

### EXAMPLE 1 : Study 1

The first study was based on a population of 109 patients, comprising patients of both nodal statuses N0 and N+. All patients were ER+ (estrogen receptor positive). All patients received Tamoxifen monotherapy immediately after surgery or diagnosis. The samples were analysed using Epigenomics' chip technology with two chip panels representing altogether 117 candidate genes. For further details see patent application WO 04/035803 and EP 03 090 432.0, which is hereby incorporated by reference. In this study one significant marker gene was found. The methylation status of PITX2, coding for a transcription factor, was correlated statistically significant with disease-free survival under adjuvant Tamoxifen treatment. A Cox regression model that includes the nodal status of the patient at the time of diagnosis was applied.

The result from this study – with respect to PITX2 - is illustrated in Figure 4. The X axis shows the metastasis free survival times of the patients in years, and the Y axis shows the proportion of metastasis free survival patients in %. Amongst the 54 patients (upper line) with below median methylation levels a higher percentage has a significantly longer metastasis free survival time, than amongst the 55 patients with above median methylation levels (lower line). To illustrate the result: At time of 10 years after surgery under tamoxifen monotherapy, more than 75% of the patients with low methylation in PITX2 are still metastasis free, whereas less than 60% of the patients with high methylation in PITX2.

As the survival of a breast cancer patient is known to also be correlated to the patient's nodal status, the differentiating power of the marker in this mixed population is expected to be less than in a homogenous population.



Another study was performed to analyse whether the same marker can be identified independently, in a completely different set of patient samples and also to characterise the differential power towards predicting survival for a sub-group of patients, all being N0.

#### EXAMPLE 2 : Study 2

The second study was based on samples from 236 patients from 5 different centres, wherein all patients were N0 (nodal status negative), and older than 35 years. In all cases the surgery was performed before 1998. All patients were ER+ (estrogen receptor positive), and the tumours were graded to be T1-3, G1-3. In this study all patients received Tamoxifen directly after surgery, and the outcome was assessed as the length of disease-free survival. In order to be as representative as possible for the final target group, the patients and their tumour samples had to fulfil the following criteria:

The range and median follow-up of patients were the following:

Median: 64.5 months

Range: 3 months to 142 months

(calculated based on patients who were disease-free at end of observation time).

Analysis of the methylation patterns of patient samples treated with Tamoxifen as an adjuvant therapy immediately following surgery (see Figure 1) is shown in the plots according to Figures 5 to 7. For the amplificate, the mean methylation over 4 oligo-pairs for that amplificate was calculated and the population split into groups according to their mean methylation values, wherein one group was composed of individuals with a methylation score higher than the median and a second group composed of individuals with a methylation score lower than the median.

The primer oligonucleotides used to generate the amplificate, that was analysed in the array experiment were these :

Array Primer PITX2\_Q21: GTAGGGGAGGGAAGTAGATGT (SEQ ID NO: 22)

Array Primer PITX2\_R23: TCCTCAACTCTACAAACCTAAAA (SEQ ID NO: 23)

The according genomic region of said amplificate is given in SEQ ID NO: 13.

---

The sequences of the oligonucleotides used in this array experiment were the following:

SEQ ID NO: NO 24 : AGTCGGGAGAGCGAAA  
SEQ ID NO: NO 25 : AGTTGGGAGAGTGAAA  
SEQ ID NO: NO 26 : AAGAGTCGGGAGTCGGA  
SEQ ID NO: NO 27: \_AAGAGTTGGGAGTTGGA  
SEQ ID NO: NO 28: GGTCGAAGAGTCGGGA  
SEQ ID NO: NO 29: GGTTGAAGAGTTGGGA  
SEQ ID NO: NO 30: ATGTTAGCGGGTCGAA  
SEQ ID NO: NO 31: TAGTGGGTTGAAGAGT

When the data derived from analysing 6 different CpG sites, located within the preferred amplified region of the PITX2 gene by means of methylation specific detection oligo hybridisation analysis were plotted as Kaplan-Meier estimated metastasis-free survival curves, it can be seen that the differential power of the marker PITX2 increased with selecting for N0 patients. This is shown in figures 5 to 7. The X axis shows the metastasis free survival times of the patients in years, and the Y axis shows the proportion of metastasis free survival patients in %. The lower curve shows the proportion of metastasis free patients in the population with above median methylation levels, and the upper curve shows the proportion of metastasis free patients in the population with below median methylation levels.

For example, as illustrated in figure 5, 10 years after surgery only about 65% of the patients of the 118 patients with the higher methylation status are metastasis free, whereas about 90% of the 118 patients with lower methylation status are metastasis free.

As illustrated in figure 6 when looking at the analogous Kaplan-Meier analysis for a subpopulation of 148 patients, characterised by a tumour at stage G1 or G2 this differential power increases again: 10 years after surgery only about 60% of the 74 patients with the higher methylation status are metastasis free, whereas about 95% of the 74 patients with lower methylation status are metastasis free.

Figure 7 illustrates how the survival is also correlated to the tumour stage at surgery by showing the analogous Kaplan-Meier analysis for a subpopulation of 150 patients, characterised by a tumour stage of T1 or T2: The number of patients with 10 years MFS is about 68% of patients of the 112 with the higher methylation status, whereas about 95% of the 112 patients with lower methylation status are metastasis free.

#### EXAMPLE 3:

The accuracy of the differentiation between the different groups was further increased by combining multiple oligonucleotides from different genes. As described in the text it was recognised that adding additional informative markers to the analysis could potentially increase the prognostic power of a survival test. Therefore it was calculated how a combination of two methylation specific oligonucleotides each from the genes TBC1D3 and CDK6, and one oligonucleotide from the gene PITX2 would differentiate the groups of good or bad prognosis. The result is shown in figure 8 as the according Kaplan-Meier curve.

Figure 9 shows -on top of Figure 8- the classification of the patients from the sample set by means of the St.Gallen method (the current method of choice for estimating disease free survival), thereby showing the improved effectiveness of methylation analysis over current methods, in particular post 80 months.

#### EXAMPLE 4: Real time Quantitative methylation analysis

Genomic DNA was analysed using the Real Time PCR technique after bisulfite conversion. In this analysis four oligonucleotides were used in each reaction. Two non methylation specific PCR primers were used to amplify a segment of the treated genomic DNA containing a methylation variable oligonucleotide probe binding site. Two oligonucleotide probes competitively hybridise to the binding site, one specific for the methylated version of the binding site, the other specific to the unmethylated version of the binding site. Accordingly, one of the probes comprises a CpG at the methylation variable position (i.e. anneals to methylated bisulfite treated sites) and the other comprises a TpG at said position (i.e. anneals to unmethylated bisulfite treated sites). Each species of probe is labelled with a 5' fluorescent reporter dye and a 3' quencher dye wherein the CpG and TpG oligonucleotides are labelled with different dyes.

The reactions are calibrated by reference to DNA standards of known methylation levels in order to quantify the levels of methylation within the sample. The DNA standards were composed of bisulfite treated phi29 amplified genomic DNA (i.e. unmethylated), and/or phi29 amplified genomic DNA treated with Sss1 methylase enzyme (thereby methylating each CpG position in the sample), which is then treated with bisulfite solution. Seven different reference

standards were used with 0%, (i.e. phi29 amplified genomic DNA only), 5%, 10%, 25%, 50%, 75% and 100% (i.e. phi29 SssI treated genomic only).

The amount of sample DNA amplified is quantified by reference to the gene ( $\beta$ -actin (ACTB)) to normalise for input DNA. For standardisation the primers and the probe for analysis of the ACTB gene lack CpG dinucleotides so that amplification is possible regardless of methylation levels. As there are no methylation variable positions, only one probe oligonucleotide is required.

The following oligonucleotides were used in the reaction to amplify the control amplicate:

Control Primer1: TGGTGATGGAGGAGGTTTAGTAAGT (SEQ ID NO: 10)

Control Primer2: AACCAATAAAACCTACTCCTCCCTTAA (SEQ ID NO: 11)

Control Probe: 6FAM-ACCACCACCCAACACACAATAACAAACACA-TAMRA or Dabcyl (SEQ ID NO: 12)

The nucleic acid sequence of the gene PITX2 is given in (SEQ ID NO: 1), after treatment with bisulfite two different strands are generated, and each of the strands is represented twice, once in a prior to treatment methylated version (SEQ ID NO: 2 and 3) and once in the prior to treatment unmethylated form (SEQ ID NO: 4 and 5), which are characterised as containing no cytosine bases (despite of those 5' adjacent to a guanine and methylated before treatment).

The following primers are used to generate an amplicate within the PITX2 sequence comprising the CpG sites of interest:

Primers for PITX bisulfite amplicate length : 144 bp

PITX2R02: GTAGGGGAGGGAAGTAGATGTT (SEQ ID NO: 6)

PITX2Q02: TTCTAATCCTCCTTTCCACAATAA (SEQ ID NO: 7)

The genomic region according to the generated amplicate of 144 bp in length is given in SEQ ID NO: NO 18.

Probes:

PITX2cg1: FAM-AGTCGGAGTCGGGAGAGCGA-Darquencher (SEQ ID NO: 8)

As an alternative quencher TAMRA was also used in additional experiments:

FAM-AGTCGGAGTCGGGAGAGCGA-TAMRA



PITX2tg1: YAKIMA YELLOW-AGTTGGAGTTGGGAGAGTGAAAGGAGA-Darquencher (SEQ ID NO: 9)

In additional experiments we also used :

VIC- AGTTGGAGTTGGGAGAGTGAAAGGAGA -TAMRA

The extent of methylation at a specific locus was determined by the following formula:

methylation rate =  $100 * I(CG) / (I(CG) + I(TG))$

(I = Intensity of the fluorescence of CG-probe or TG-probe)

PCR components were ordered from Eurogentec :

3 mM MgCl<sub>2</sub> buffer, 10x buffer, Hotstart TAQ

Program (45 cycles): 95 °C, 10 min; 95 °C, 15 sec; 62 °C, 1 min

This assay was performed on 236 samples identical to those used in Example 2. The result is shown in figure 2. Figure 2 shows the Kaplan-Meier estimated disease-free survival curves for 3 CpG positions of the PITX2 gene by means of Real-Time (RT) methylation specific probe analysis, as described above. The lower curve shows the proportion of disease free patients in the population with above median methylation levels, the upper curve shows the proportion of disease free patients in the population with below median methylation levels. The X axis shows the disease free survival times of the patients in months, and the Y- axis shows the proportion of disease free survival patients. The p-value (probability that the observed distribution occurred by chance) was calculated as 0.0031, thereby confirming the data obtained by means of array analysis.

For comparison, figure 3 illustrates the result from the array analysis of said gene, according to the chip hybridisation experiment described in Example 2, wherein detection oligos were used (for details see EP 03 090 432.0, which is incorporated by reference). The p-value (probability that the observed distribution occurred by chance) was calculated as 0.0011.

#### EXAMPLE 5

Another QM assay was developed in our hands, which also performed very well. The following PITX2 specific oligonucleotides were employed to generate an amplicate of 164 bp

The oligonucleotides are specific for three co-methylated CpG positions:

Primers for PITX2 bisulfite amplificate with a length of 162 bp :

PITX2O2: AACATCTACTTCCCTCCCCTAC (SEQ ID NO: 14)

PITX2P3: GTTAGTAGAGATTTTATTAAATTTTATTGTAT (SEQ ID NO: 15)

The genomic region according to the generated amplificate of 162 bp in length is given in SEQ ID NO: NO 19.

Probes (from ABI):

PITX2-IIcg1: FAM-TTCGGTTGCGCGGT-MGBNQF (SEQ ID NO: 16)

PITX2-IItg1: VIC-TTTGGTTGTGTGGTTG- MGBNQF (SEQ ID NO: 17)

The extent of methylation at a specific locus was determined by the following formula:

methylation rate =  $100 * I(CG) / (I(CG) + I(TG))$

(I = Intensity of the fluorescence of CG-probe or TG-probe)

PCR components were ordered from Eurogentec : 2,5 mM MgCl<sub>2</sub> buffer, 10x buffer, Hotstart TAQ

Program (45 cycles): 95 °C, 10 min; 95 °C, 15 sec; 60 °C, 1 min

### Example 6

#### Patient material

The material to be used in this study; consists of fresh frozen healthy breast tissue, fresh frozen breast tumour tissue from untreated breast cancer patients (follow up over >10 years) and samples from Tamoxifen treated patients (follow up over >10 years from Tamoxifen treatment). Aliquots of DNA from these microdissected lesions are used as the source template for PCR-based LOH (Loss of heterozygosity) analysis. All tumour samples were derived from ER+ node negative patients.

#### LOH analysis

DNA from all tissue samples is subjected to PCR-based LOH analysis using two 4q25-26 markers (D4S1284 and D4S406). These markers define a region on chromosome 4 comprising the gene PITX2 gene said region but being more than 8.5 kbp distant of a region previously shown to undergo LOH in breast carcinomas [*Cancer Research* 59, 3576-3580, August 1, 1999].

### DNA Extraction

Extract DNA from samples using the Wizzard Kit (Promega).

### PCR reaction

See Clin. Cancer Res., 5: 17-23, 1999 for further details.

Analyse each sample by means of single-plex PCR using the following primers:

D4S406

Forward primer: GAAAGGCAGAGTCATAACAGGAAG (SEQ ID NO: 32)

Reverse primer: TAAGGATAGAGTGATTTCCAAGAAAG (SEQ ID NO: 33)

PCR product size: 205 (bp)

GenBank Accession: Z16728

D4S1284

Forward primer: CTTATCTGACAACAAGCGAGTATG (SEQ ID NO: 34)

Reverse primer: CAATTATTGTATTGTAGCATCGGAG (SEQ ID NO: 35)

PCR product size: 172 (bp)

GenBank Accession: L14168

Synthesise forward primers with either a fluorescent FAM tag (D4S1284) or a fluorescent TET tag (D4S406) at the 5' end.

Prepare a suitable quantity of nucleotide mixture according to Table 2.

Aliquot 1 µl of each DNA sample into separate PCR tubes, add 9 µl reaction mixture according to Table 3 and thermal cycle according to the following conditions.

Thermal cycling conditions:

95°C for 15 min

39 cycles:

95°C for 1 min

55°C for 0:45

72°C for 1:15

72°C for 10 min

---

### Gel electrophoresis

Horizontal ultrathin, high throughput fluorescence-based DNA fragment gel electrophoresis is the preferred technique to separate and analyse the PCR-generated alleles. Combine one microliter of amplified material with 2  $\mu$ l formamide loading dye (APB) prior to electrophoresis. Add ROX 350 fluorescent size markers (0.7  $\mu$ l; ABI) to amplified tumour DNA to allow sizing of alleles.

Heat samples to 95°C, load on 70  $\mu$ m, 5% horizontal polyacrylamide gel and electrophorese for 1 h and 15 min at 30 W in 1  $\times$  TBE.

Data may be collected as commonly known in the art (see for example Clin. Cancer Res., 5: 17-23, 1999). To determine whether allelic deletion had occurred at individual markers, calculate allelic ratios and express as a percentage of loss of intensity for the treated and untreated tumour samples compared with the corresponding normal samples (D-value) after normalisation. When the allelic ratio in the tumour DNA is reduced by greater than 40% (DO.40) from that found in the normal DNA, the sample is denoted as having LOH at that locus.

Table 2: Nucleotide Mix

10 $\mu$ l dATP, 10 mM
10 $\mu$ l dGTP, 10 mM
10 $\mu$ l dTTP, 10 mM
2.0 $\mu$ l dCTP, 10 mM
288 $\mu$ l DEPC-treated H <sub>2</sub> O

Table 3: Reaction mixture

1.0 $\mu$ l Taq Buffer
0.8 $\mu$ l Reduced nucleotide mixture
0.2 $\mu$ l Forward primer, 20 $\mu$ M
0.2 $\mu$ l Reverse primer, 20 $\mu$ M
6.6 $\mu$ l DEPC treated H <sub>2</sub> O
0.1 $\mu$ l $\gamma$ -32P dCTP
0.1 $\mu$ l AmpliTaq Gold Poly- merase



Total volume = 9  $\mu$ l

16. Nov. 2004

## CLAIMS

1. A method for characterising a cell proliferative disorder of the breast tissues and/or predicting the survival of a patient diagnosed with said disorder, comprising the steps of:
  - (a) detecting the expression of a nucleic acid or a polypeptide expressed from the PITX2 gene in an isolated biological sample representative of said cell proliferative disorders of the breast tissues and
  - (b) therefrom predicting the survival of said patient, characteristics of said cell proliferative disorder, and/or prognosis of said patient.
2. The method according to claim 1 further comprising
  - (c) determining a suitable treatment regimen for the subject.
3. The method of claim 1, wherein said patient is characterised by being subject to adjuvant endocrine therapy comprising one or more treatments which target the estrogen receptor pathway or are involved in estrogen metabolism, production or secretion.
4. The method of claim 1, wherein said breast cell proliferative disorders are taken from the group comprising ductal carcinoma *in situ*, invasive ductal carcinoma, invasive lobular carcinoma, lobular carcinoma *in situ*, comedocarcinoma, inflammatory carcinoma, mucinous carcinoma, scirrhous carcinoma, colloid carcinoma, tubular carcinoma, medullary carcinoma, metaplastic carcinoma, and papillary carcinoma and papillary carcinoma *in situ*, undifferentiated or anaplastic carcinoma and Paget's disease of the breast.
5. The method according to claim 1 characterised in that the detection is carried out by
  - a) contacting said biological sample with an antibody immunoreactive with the PITX2 polypeptide to form an immunocomplex;
  - b) detecting said immunocomplex; and
  - c) predicting therefrom the survival of said patient, characteristics of said cell proliferative disorder, and/or prognosis of said patient.
6. The method according to claim 1 characterised in that the detection is carried out by

- a) contacting said biological sample with an antibody immunoreactive with the PITX2 polypeptide to form an immunocomplex;
- b) detecting said immunocomplex;
- c) therefrom predicting the survival of said patient, characteristics of said cell proliferative disorder, and/or prognosis of said patient, and
- d) comparing the quantity of said immunocomplex to the quantity of immunocomplex formed under identical conditions with the same antibody and a control sample from one or more patients with a known prognosis.

7. The method according to claim 1 characterised in that the detection is carried out by

- a) contacting said biological sample with an antibody immunoreactive with the PITX2 polypeptide to form an immunocomplex;
  - b) detecting said immunocomplex; and
  - c) therefrom predicting the survival of said patient, characteristics of said cell proliferative disorder, and/or prognosis of said patient,
- and wherein an increase in quantity of said immunocomplex in the sample from said subject relative to said control sample is indicative of a bad prognosis.

8. The method of claim 5, wherein said immunoassay is a radioimmunoassay or an ELISA or a Western blot.

9. The method of claim 1, wherein said detection is afforded by mRNA expression analysis.

10. The method of claim 9, comprising detecting the level of mRNA encoding a PITX2 polypeptide in a biological sample from a patient,

11. A method according to claim 10, wherein a increased concentration of said mRNA above the concentration determined for an individual known to have a good prognosis indicates a bad prognosis.

12. The method of claim 10, comprising the steps of:

- (a) providing a polynucleotide probe which specifically hybridises or is identical to a polynucleotide consisting of SEQ ID NO: 19 or SEQ ID NO: 1,

- (b) incubating said sample with said polynucleotide probe under high stringency conditions to form a specific hybridisation complex between a nucleic acid and said probe;
- (c) detecting said hybridisation complex.

13. The method according to claim 12 wherein said nucleic acid is mRNA or a cDNA derived therefrom.

14. The method according to claim 13 wherein the detecting step further comprises the steps of:

- a) producing a cDNA from mRNA in the sample;
- b) providing two oligonucleotides which specifically hybridise to regions flanking a segment of the cDNA;
- c) performing a polymerase chain reaction on the cDNA of step a) using the oligonucleotides of step b) as primers to amplify the cDNA segment; and
- d) detecting the amplified cDNA segment.

15. Use of a polypeptide expressed from the PITX2 gene for differentiating or distinguishing between patients diagnosed with breast cancer, who have a good survival prognosis and patients who have a bad survival prognosis.

16. Use of a polypeptide expressed from the PITX2 gene for prediction of survival of a patient diagnosed with a cell proliferative disorder of the breast.

17. The method of claim 1 wherein said detection comprises determining the genetic parameters of the gene PITX2, its promoter and/or regulatory elements.

18. The method of claim 1 wherein said detection comprises determining the epigenetic parameters of the gene PITX2, its promoter and/or regulatory elements.

19. The method of claim 1, wherein said detection comprises determining the methylation status of one or more CpG positions of a target nucleic acid within the gene PITX2, its promoter and/or regulatory elements, in particular through the methylation analysis of a genomic DNA sequence according to SEQ ID NO: 1.



20. The method of claim 1, wherein said detection comprises determining the methylation status of one or more CpG positions of a target nucleic acid characterised as being identical to or hybridising under stringent or moderately stringent conditions to a sequence out of the group of SEQ ID NO: NOs 13, 18 and 19.

21. The method of claim 19, wherein the methylation analysis is afforded by contacting said target nucleic acid with one or more agents that convert cytosine bases that are unmethylated at the 5'-position thereof to a base that is detectably dissimilar to cytosine in terms of hybridisation properties.

22. The method of claim 21, wherein contacting said target nucleic acids with one or more agents comprises use of a solution selected from the group consisting of bisulfite, hydrogen sulfite, disulfite, and combinations thereof.

23. Use of a set of oligomer probes comprising at least two oligomers, in particular an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to a pretreated genomic DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for detecting the cytosine methylation state and/or single nucleotide polymorphisms (SNPs) within one of the sequences according to SEQ ID NO: 1, and sequences complementary thereto.

24. A method for manufacturing an arrangement of different oligomers (array) fixed to a carrier material for analysing diseases associated with the methylation state of the CpG dinucleotides of one of SEQ ID NO: 1, and sequences complementary thereto wherein at least one oligomer, in particular an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to a pretreated genomic DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, is coupled to a solid phase.

25. A composition of matter comprising the following:

- a nucleic acid comprising a sequence at least 18 bases in length of a segment of the

chemically pretreated genomic DNA according to one of the sequences taken from the group comprising SEQ ID NO: 1 to SEQ ID NO: 5 and sequences complementary thereto, and

- a buffer comprising at least one of the following substances: 1 to 5 mM Magnesium Chloride, 100-500  $\mu$ M dNTP, 0.5-5 units/10ul of taq polymerase, an oligomer, in particular an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to a pretreated genomic DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto.

26. Use of the gene PITX2, its promoter and/or regulatory elements for predicting the survival of patients diagnosed with a cell proliferative disease.

27. Use of the mRNA of the gene PITX2 for predicting the survival of patients diagnosed with a cell proliferative disease.

28. A method for predicting the survival of patients diagnosed with a cell proliferative disease according to claim 19, comprising:

- a) isolating or enriching genomic DNA from said biological sample ;
- b) treating the genomic DNA, or a fragment thereof, with one or more reagents to convert 5-position unmethylated cytosine bases to uracil or to another base that is detectably dissimilar to cytosine in terms of hybridisation properties;
- c) contacting the treated genomic DNA, or the treated fragment thereof, with an amplification enzyme and at least two primers comprising, in each case a contiguous sequence at least 18 nucleotides in length that is complementary to, or hybridises under moderately stringent or stringent conditions to a sequence selected from the group consisting of SEQ ID NO: 2 to 5, and complements thereof, wherein the treated DNA or a fragment thereof is either amplified to produce one or more amplicates, or is not amplified; and
- d) determining, based on the presence or absence of, or on the quantity or on a property of said amplicate, the methylation state of at least one CpG dinucleotide sequence of SEQ ID NO: 1, or an average, or a value reflecting an average methylation state of a plurality of CpG dinucleotide sequences of SEQ ID NO: 1.

29. A method for detecting the survival of patients diagnosed with a cell proliferative disease of the breast according to claim 19, comprising the following steps of

- a) obtaining, from a subject, a biological sample having subject genomic DNA;
- b) treating the genomic DNA, or a fragment thereof, with one or more reagents to convert 5-position unmethylated cytosine bases to uracil or to another base that is detectably dissimilar to cytosine in terms of hybridisation properties;
- c) amplifying one or more fragments of the treated DNA such that only DNA originating from breast or breast cell proliferative disorder cells are amplified
- d) detecting the amplificates or characteristics thereof and thereby deducing on the survival of patients diagnosed with a cell proliferative disease of the breast.

30. Use of an oligomer, an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to an artificially modified, chemically pretreated DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for differentiating or distinguishing between patients diagnosed with breast cancer, who have a good survival prognosis and patients who have a bad survival prognosis.

31. Use of a nucleic acid comprising a sequence of at least 18 bases in length of a segment of the artificially modified, chemically pretreated, DNA according to one of the sequences taken from the group comprising SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for differentiating or distinguishing between patients diagnosed with breast cancer, who have a good survival prognosis and patients who have a bad survival prognosis.

32. Use of an oligomer, an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to an artificially modified, chemically pretreated DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for prediction of survival of a patient diagnosed with a cell proliferative disorder of the breast.

33. Use of a nucleic acid comprising a sequence of at least 18 bases in length of a segment of the artificially modified, chemically pretreated, DNA according to one of the sequences taken from the group comprising SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for prediction of survival of a patient diagnosed with a cell proliferative disorder of the breast.

34. Use of a nucleic acid represented by their SEQ ID NO: NO out of the group of nucleic acids according to SEQ ID NO: NOS : 6, 7, 8, 9, 10, 14, 15, 22, 23, 24, 25, 26, 27, 28, 29, 30 and 31, for differentiating or distinguishing between patients diagnosed with breast cancer, who have a good survival prognosis and patients who have a bad survival prognosis.

35. Use of a nucleic acid represented by their SEQ ID NO: NO out of the group of nucleic acids according to SEQ ID NO: NOS : 6, 7, 8, 9, 10, 14, 15, 22, 23, 24, 25, 26, 27, 28, 29, 30 and 31, for prediction of survival of a patient diagnosed with a cell proliferative disorder of the breast.





**Abstract**

**16. Nov. 2004**

The present invention relates to methods for predicting the survival of a human being diagnosed with a cell proliferative disorder of the breast tissues, characterised by a step of determining the expression level of PITX2 or the genetic or the epigenetic modifications of the genomic DNA associated with the gene PITX2. The invention also relates to sequences, oligonucleotides and antibodies which can be used within the described methods.



Figure 1






Figure 2




Figure 3




Figure 4




Figure 5






Figure 6




Figure 7




Figure 8




Figure 9






Figure 10

METNCRKLVSACVQLGVQPAAVECLFSKDSEIKKVEFTDSPESRKEAASSKFFPRQ  
HPGANEKDKSQQGNEDVGAEDPSKKRQRRQRTHTSQQLQELEATFQRNRYYP  
DMSTREEIAVWTNLTEARVRVWFKNRRRAKWRKRERNQQAELOCKNGFGPQFNGL  
MQPYDDMYPGYSYNNWAAKGLTSASLSTKSFPFFNSMNVNPLSSQSMFSPPNSSISS  
MSMSSSMVPSAVTGVPGSSLNSLNNLNNLSSPSLNSAVPTPACPYAPPTPPYVYRDT  
CNSSLASRLKAKQHSSFGYASVQNPASNLSACQYAVDRPV

Sequence listing

16. Nov. 2004

<110> Epigenomics AG

<120> PITX2 - a marker to predict survival of patients diagnosed with  
breast cell proliferative disease

<160> 35

<210> 1

<211> 9001

<212> DNA

<213> Homo Sapiens

<400> 1

agctgatgga	cttgctaaat	ttctttcttc	ttttttcttt	ttcatattat	ttgctagcca	60
taatggaatc	ctctagggtt	aagccaaaga	aaaattggag	agacaaaatt	agattttgta	120
gcccttttcc	cccccgga	tgcctttttt	tttcttttta	gtttctgatg	aatggctatc	180
atttatttct	accaaattta	aataaggact	gctgccttgt	atgtttaact	aggcaggcag	240
agggaactgg	tttgtttagg	aagcagtgc	tgagatgtcc	tggccaagtt	agtgacagag	300
gaggggagaa	agaatccaga	ccaatttgta	tgcagtatat	tttactccca	tgaaataaaa	360
cacatttggt	tcatatttgc	tgaaaagtaa	aacaataata	ttgtacgaaa	tggtatacac	420
agggtagggt	gtacatagca	gtttcagaaa	catcattgca	tccaccagag	aaactattct	480
aaaactgata	ttcacacatt	ttttataata	ataataatat	gttagaaaca	tacagtgtgg	540
catttagtat	atacactccc	ttgctcgcaa	gcgaaaaatc	ctaactcgctt	ctgtataaca	600
tgcttttatt	taaagcctaa	ccttttaaaa	cactgttggt	atattactaa	caactgcttt	660
tataaaatta	atttgacatt	tcgatataata	tacatccttt	cagtcattta	aatgttaaca	720
atgctaaact	taaaaaataa	caagcttata	gtaatgttaa	aatgtcatat	ccagtcaaac	780
atttgtttgt	gtatgtgtcc	ttgcaactgt	tagaaataact	tgtagtgaag	gatgtcagac	840
actgaggaca	tccctttgaa	atcaaaggag	ctctctcttt	gattcagtgg	tttcttttct	900
tctatatagc	ttctctttct	ctccctttct	ttagtgccca	cgaccttcta	gcataattcc	960
cagtctttca	agggcggagt	tgcccatcc	ggcaagggtcc	taggatcccg	gcgctgtggg	1020
tgcggtcac	acgggcoggt	ccactgcata	ctggcaagca	ctcagggttg	aggccgggtt	1080
ctgcacgctg	gcgtagccga	agctggagtg	ctgctttgct	ttcagtcctca	ggctggccag	1140
gctcgagtta	cacgtgtccc	tataaacata	cggaggagtc	ggcggcgcgt	aaggacaggc	1200
aggcgtcggc	accgcggaat	tcagcgacgg	gctactcagg	ttgttcaagt	tattcaggct	1260
gttgagactg	gagcccggga	cgctgtcac	tgctgagggc	accatgctgg	acgacatgct	1320
catggacgag	atagagttgg	gtggggaaaa	catgctctgt	gatgacaggg	ggttgacggt	1380
catagagttg	aagaaggggga	agctcttggt	ggatagggag	gcggatgtaa	ggcccttggc	1440
ggcccagttg	ttgtaggaat	agcctgggta	catgtcgtcg	tagggctgca	tgagcccatt	1500
gaactgcggc	ccgaagccat	tcttgcatag	ctcggcctgc	tgggttgcgt	ccctctttct	1560
ccatttgggc	cgacgattct	tgaaccaaac	ctggggggcg	ttggggcaag	ggagcaaaca	1620
gatgccacag	tgcagattac	taaaacttcc	atcgaggagg	aacccccgcc	ttcccccgac	1680
acacacgcta	gcgcactcac	acaccctggc	ctcgcttcac	tgcaccgccc	tgacaccaa	1740
gataccaggg	ccagctttca	gttactggcc	cgggtctcca	ccaagcgag	gagacctggt	1800
ctgctctggc	ctgcgagctg	ggactcggag	ctacgccaca	aacctcagcc	gaacgcatgg	1860
agacctgcgg	acggtttgat	cactcagcca	ggcgtttctc	cagggtccaa	aacacttaat	1920
gtaaaacaaa	cgcgggggcag	caggcttttc	caacccttcc	cggggcacct	tgcaaacttg	1980
cttccattcc	aaagccacag	acccacggat	gaggagaagg	ggctggaagg	gcactagagg	2040
atcgctcttt	ctcccacgca	attcctccct	tccttccctg	acctccactg	tcgtcccca	2100
ccccctggta	cgtgctccct	taacagggac	taggcgcgca	acactctttc	tcgcctagca	2160
aaacaaccaa	ataaagagca	aaagaccacc	tcttcgtcag	ctcgtttaact	ccaggagctt	2220
ggcatattaa	actccgggaa	cccggaaagg	gtagttttgg	agattccccc	ttctttcgct	2280
ctgcctcttc	tttaccctaa	gcccaccaca	ggcctgtccg	cgcgccaggc	ccagccgggt	2340
cgtttggtt	tgcaggcggc	caccagggc	ggcgggttc	caccggtgtc	cggtggcca	2400
gccgcaacc	cgatcccaat	ccacatcggt	cctccctgtc	gccccagacg	gcggcttttg	2460
tgtattggag	agaggcctgg	cctgagatat	ccgagctgac	accagtgatg	tttcacatta	2520
cacatctccg	ccggggcccag	ccgtgtaate	cgctttttct	ctttttcctt	tcattcttga	2580
tttctttttt	atcccccttc	ctctttgcac	ccgactgcta	taaaaagcac	gcctcactcc	2640
cacttggttc	gacaagcagc	cgccctggaa	ggagaggcag	ctgcaaggag	agcccagcgc	2700
cgcggttaca	aagcactagg	gtggagctgc	ggaatagcgg	gcgggggtgg	agggcgtttt	2760
cgaaggatcc	cagaaaaccc	atagactctg	tctttaatta	cttgccattt	ctaccctagg	2820

ccatctaatac	tttgctcagg	cgagaagagt	acgtgagagg	cccgttccct	tgatgtgcaa	2880
gagagctaata	gaaagactga	ccttgctcaa	aaccacgccg	cccaggaccc	agctctggct	2940
ctggacagtt	aaactaaaac	cattttcaac	ttcttcccgg	ccttttatcc	accagcatag	3000
cctcatgcct	tgcacaaatg	ccacccagag	agtgtcttca	ttccctctga	tttgggagag	3060
cattttggct	tttattcttt	ttatcgttgt	tttcttcttt	ttgtttgctc	tgctctaacc	3120
gggggcttta	ttttttctac	ccagagcact	taattttttt	tttttaacag	caaagcctct	3180
ggatgccgct	tgatttgctt	gattctgttt	tctgcttcca	gaatcctaac	aaatttgga	3240
tcttccaccg	accagcataa	accaggacgt	tgctattggg	ttattttatt	gagctcattt	3300
ttgccaatcc	ataaagtaca	gatttgctac	aaagttaagg	taagcccttt	ttacaaaact	3360
atgattataa	tttagaagag	gggtgtgag	tttcaatttc	cagagttcaa	ctcctgagag	3420
aagataaata	aaccaagcag	aaaagtcttt	cttctttttt	tctttctcct	tctaagagga	3480
ctagtagttg	tgtattaaaa	ctttgctccc	ggagatcaca	aaactaggaa	ataggggtgt	3540
tgggagagac	ctgaatggcc	gaaacaaccg	taaagaaggt	gtaagaagcg	cgagcccagg	3600
agggaaaaag	ctgggccagg	gccgggacaa	aggtttccca	gggagggcca	actcttccgt	3660
gtctctggcg	ggttttcctt	gttaaaggct	cacagggttg	agcctgttcg	cggctcttgg	3720
cctggtaggg	atthttattag	ctctgctctg	gcaactgcaa	gccaggaaca	caatgtcctg	3780
tgcaggggat	tgcccatgca	gccagctcgc	tgagatcgcg	ggatggcggg	gcagtgagcc	3840
ggtgccgctc	tgggagcctg	agccaggcg	gcagtcctgt	cggcctcgga	gagggaaactg	3900
taatctcgca	accaggccgc	cgcgaggcct	tctgcctttg	caaagctgcg	ccccaccggc	3960
gcctcccag	gcggcgctgc	cttccacatt	ctctcctggg	ctacttgggc	tgtacctcca	4020
caacatcctc	cccccatccc	tcccagactc	cgtgctggct	cctacccgga	ctcgggcttc	4080
cgtaagggtg	gtccacacag	cgatttcttc	gcgtgtggac	atgtccgggt	agcgggtcct	4140
ctggaaagtg	gcctccagct	cctggagctg	ctggctggta	aagtgaagtc	gctgccgcct	4200
ttgccgcttc	ttcttagacg	ggtcctcggc	gccacgctcc	tcattcttcc	cctgctggct	4260
tttatctttc	tctgaaaacg	aaacacacac	actttcccgt	cagcatgccc	acctgcaacg	4320
cggacgcca	ctggaccggc	ggcagaagcc	gtggaagagc	tgggctgcct	ggcgccggag	4380
gaggggtgcg	gcggcggtc	cgggcccgcg	ggagcgctgc	gcctgtgggg	tgtgcaggcg	4440
caagtgtggg	tgtccgcgc	ccatttcttc	ccctccccca	gcgcgcgacg	ttttatttac	4500
atgtttatct	cactgcagcg	gcacattcac	ttttatagcc	tgtgctttca	agtatatatta	4560
tacacctctg	cgcagacaca	ccaaatctcc	tgggacgcgc	acacgcgcgt	ggtttacaga	4620
ccccctccc	cctcgcagaa	agctcagatt	tccatgcggt	ttgggaaggc	taggaaaaga	4680
tgtggggatt	cgggttgggca	ccgaagtctg	ccggcccttt	cccaaaaaaa	aaaaaaaaaat	4740
gcctcttcgc	gaagggcatt	tctgagtggg	ttcaggcaat	ttcctaacga	gtggagctcc	4800
tccggagctg	aaagccgaga	ggaaaacagg	gacagaggtc	ggcggcctct	gaaggctctc	4860
gaatcaagat	gctgggattt	ttgtgaccca	ggaaacagaa	gggaggccag	ggtacgaata	4920
gagagggcg	cagaattgct	cgcgcctta	gcgcccagag	agccggggcg	gtcgagggag	4980
aactaaagg	atgcggggta	gtcaaaattc	cggctcccgg	aagtctcgcg	gggagccagg	5040
cgaacgacca	ctcccaccac	gcctcccccc	ggaggggctg	acttcccttg	ggcgagaggg	5100
agcgggtggc	gcagagcagc	tgagcgggaa	tgtctgcagg	gcggcgcggc	gccttacctg	5160
cggcctccgg	gctggaggtg	tccggagatg	tgtgcacctc	cagcctgtgc	ttggaggagt	5220
ccagcgaccg	gggctgaccg	ggagccagaa	ccgaagccat	ggctaaccgc	tggggatggt	5280
gacaggaaga	tgaggagacg	gccgacagct	tggtccccgc	tgtcgggtgc	tccaagtga	5340
gcgggccttt	catgcagttc	atggacgagg	gagcgcgacg	ctctactagt	ccttggctac	5400
tgccccgcgg	agcccccgta	gccgcgcgtg	cccgcctccg	gtcgcgctct	aggcgcggag	5460
tttccccgct	gcgggggagag	ccagggggacg	caacccccgc	cgagttctca	agccaagctg	5520
cccccgcttc	ctccggaagg	ctcaagcgaa	aaagtccgga	gacggaaagt	cagcgggcaa	5580
acgaagacat	gggatgtggg	cagaagggca	ccactcagag	cgtcttttag	gagcaggctt	5640
ccaagctcca	aagcgaaaca	agagtgggca	aagacccctt	tcttctctcc	ctccctcccc	5700
caagaacccc	tccaataagg	aaagctaacg	ccgaccgcgc	tctgcccgc	ccccccccac	5760
gcggcagccc	tgacagagaa	gtgtcaagag	tgacagggac	aggtaggtga	tattagatcc	5820
cctgcggcg	cagcagccgc	tgcagccacg	acgcggccct	ctgagcgcac	cctccgcaac	5880
gcgcacacgc	acacccctcg	ggcggtcgaa	caggagccgg	gccttgccgc	agctcagctc	5940
caggcaccca	ggcgagcgac	ggaccagatc	tgcggctccg	cgttccctcg	ttggcctaac	6000
atcttaaaac	cagaggcggg	cttccctggg	ccgagacgtc	actccgcgc	ggccctcccc	6060
agccctctcc	gcctccgcct	cctcccagac	ccttctccgg	gtgcgactga	cgtggctccg	6120
caccaatcag	gacgccccga	gccgcggtgg	agggactgtc	ctgcctgcac	ctatcagcag	6180
tgcggggcg	ggctactgcc	tgcgcgtg	cactgggtct	acacaggcaa	gctcccggga	6240
attcagctcc	tgcccagccc	aaggcgatcc	ggcttttagt	acgaacccaa	agggtgaagag	6300
atgaggctag	gagtcgaagg	cttgggagaa	gagagtggaa	tggtaagaa	gagaaaggta	6360
caaggatcaa	caagacaccc	actctttgtg	tctcactaca	tccattttcca	atccccacc	6420
ccatataaaa	aggagacacg	ttacttaaaa	ctagaaaatt	tgaaaaaacag	caacaaatca	6480
cctctccgat	cttaaattht	ccaaacagcc	tgtcaagtga	atgctgcgct	aatctgaaga	6540
agctttaatt	gcaaagaaga	cagagccctg	aaaaggcagg	ctaataaatt	agaaatcgag	6600



aagcaaatgg	acccgtcaaa	agaaaattac	cttgacttta	aacgaacaac	tgttttggtgg	6660
ttcactctgg	at ttatacaa	gaataaaaag	tgcctcaga	tcacgttctc	tgtgatgctt	6720
attagtcctc	agacagaaaa	cacacaatag	aagagaaacc	ctaaccacag	gtttttcaaaa	6780
tgctgaaagc	ttatccattc	tacttaacgt	tgattaagac	acatatccta	gatcttttcaa	6840
attccttgta	cactgtatta	agctcgtcct	aaccogagag	agccacgctt	taaattcgac	6900
tctcttggtt	actttattat	caatcagatt	taaatccata	aagcctgtag	aatcaacaac	6960
cttgagctaa	ttatatatga	aatatgcctt	aatgaatttc	catacaatta	agaatggtgc	7020
caaataacca	at ttcaagga	taatttttta	cagtcatttt	cttttcccag	tgagctcaag	7080
gctgtcttga	gccattaaag	tccaagcagg	cagaaggggt	gtgtgtgagc	taagggcgaa	7140
aagcctagaa	ctgcgctcaa	ctagcaaaaag	caaaacctta	tttatataaa	acaaaaaaa	7200
tcacctttgg	agacatcaac	tctttatagc	actgtttcca	agcaaattta	at ttccaaag	7260
aaattaaaga	aagaaatcca	aacatattca	aaataatttt	tgaaagtcct	tttgtccccc	7320
agcataggtc	agctggagag	gacaaactaa	tctcctctgg	gtttctgcat	gggcgattgt	7380
tttactatgg	agttagtgtt	atcatctctg	aatgtgtatt	tgtttgacat	tacagtcaat	7440
gatttgcaat	gccagcatga	agtatcttta	aaacactccc	tccttgtcct	tgttcacaaag	7500
attgggaaac	ttattcgatg	tggaacaaag	tggatgaagc	agactacaaa	tatat ttgca	7560
acttatgtgt	ctctcctttg	ccctgaccac	cccaaacc	tatctgcaac	tcctcccat	7620
tttaaaactg	cagtccaaag	acgcacatga	gaattgtttt	tcagtccttc	ttcaccagta	7680
tcacccact	ttaagaataa	tttagctgca	agggaggaat	ttcttcatag	taagctttaa	7740
atcagcattt	ctgcttttta	ccttttattc	cactttaccc	cattccacac	atacagacac	7800
ctgctcagag	taaaacacat	cctcatgtga	caggtctgca	ttagctgagg	ctcatacatc	7860
cagctatatt	aggtcctgca	atccttatcac	taaattatac	acattacact	agcagcctgt	7920
tggtaaagaa	ggttaaatta	at ttacattc	tgctcattat	ctgggtgctta	aatgacgcat	7980
tttatcccgg	agatttggcg	gagaatctcc	ttctcagacc	ccacagcgtt	tcactgaaga	8040
caatgctctt	acatttgtag	tggtttttta	tctgataaga	ctctaatttg	cttaagtctt	8100
ttaaataagg	gttttaaatg	tttctagccg	ttttcttatt	gaatttcctc	taattccccc	8160
aagatcataa	agtatatgtg	taaagtaa	at ttccctcc	attgcaactgc	cagccgatga	8220
cctataacta	agtcaataag	aatccagctc	ttttctgctg	aatgtgttta	ctaatacat	8280
tccagtttct	tcttttaaac	ctcagaatag	ctgtgggtccc	cacaatacca	tgccccttaa	8340
agcttcattt	catgaaggga	ctccatcaca	ttaaagaatg	aaaaaaatct	ccactgtagt	8400
tagtatacac	agccctcac	tccttggttt	tcaagattca	aacccacagag	ctgcaaatat	8460
ttttggaagc	ttgggtgtta	atgccttatt	ttagaaagcc	gagaagcccc	acagagccat	8520
atagatttct	aaacccatct	ctcataaacc	cacagaat tt	tgataaaaagc	tctgggtggct	8580
ccactctacc	gatggaactt	tcacacagac	aaatatacat	gtatgaagga	cctcaatcag	8640
cctccaaagt	ggttgaaaaa	cccaagggca	cgtgactgct	cctcatagtg	ccaacgtgtg	8700
cgagatgttg	gaagcactgg	ggatcagcag	cagcctagat	gcctaaaaag	ataaggtgtc	8760
ctaatttggtg	tggacccatt	gaagtcaagt	ggtgaataaa	gacaattatc	tagataattc	8820
agattaaagt	aaaagcaaaa	ccatatctat	ttgtatatat	atattcacat	ccattttata	8880
ttacagacac	acacacgcac	acacacactg	gctctgtaaa	caactgactc	aaagtgagga	8940
ttttctttgc	at ttttccag	caggagtttc	aacattctcc	taatctccta	atcactttac	9000
a						9001

<210> 2

<211> 9001

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 2

agttgatgga	tttgttaaat	tttttttttt	tttttttttt	tttatattat	ttgttagtta	60
taatggaatt	ttttagggtt	aagttaaaga	aaaattggag	agataaaatt	agatttttga	120
gttttttttt	ttttcgggaa	tggttttttt	ttttttttta	gttttttgatg	aatgggttatt	180
at ttat tttt	attaaattta	aataaggatt	gttggtttgt	atgttttaatt	aggtaggtag	240
agggaattgg	tttgtttagg	aagtagtgat	tgagatgttt	tgggttaagtt	agtgatagag	300
gaggggagaa	agaatttaga	ttaatttgta	tgtagtatat	tttat ttta	tgaaataaaa	360
tatat ttgtt	ttatat ttgt	tgaaaagtaa	aataataata	ttgtacgaaa	tgttatatat	420
agggtagggt	gtatatagta	gttttagaaa	tattatttgta	tttattagag	aaattatttt	480
aaaattgata	tttatatat	ttttataata	ataataatat	gttagaaata	tatagtgtgg	540
tatttagtat	atatat tttt	ttgttcgtaa	gcgaaaaatt	ttaatcgttt	ttgtataata	600
tgttttat tt	taaagtttta	tttttaaaaa	tattgttggtg	atattattaa	taattgtttt	660



tataaaatta	atattgatatt	tcgatatata	tatatTTTTT	tagttattta	aatgttaata	720
atgttaaatt	taaaaaataa	taagtttata	gtaatgttaa	aatgttatat	ttagttaaat	780
atattgtttgt	gtatgtgttt	ttgtaattgt	tagaaatatt	tgtagtgaat	gatgttagat	840
attgaggata	TTTTTTTgaa	attaaaggag	TTTTTTTTTT	gatttagtgg	TTTTTTTTTT	900
tttatatagt	TTTTTTTTTT	TTTTTTTTTT	ttagtggtta	cgatttttta	gtataatttt	960
tagtttttta	agggcggagt	tgTTTTtattc	ggtaagggtt	taggatttcg	gcgttgtggg	1020
tgcggtttat	acgggtcggg	ttattgtata	ttggtaagta	tttaggttgg	aggtcgggtt	1080
ttgtacgttg	gcgtagtoga	agttggagtg	ttgttttggt	tttaggttta	gggtggttag	1140
gttcgagttt	tacgtgtttt	tataaatata	cggaggagtc	ggcggcgcgt	aaggataggt	1200
agggcgcggg	atcgcggaat	ttagcgacgg	gttatttagg	ttgtttaagt	tatttaggtt	1260
ggttgagattg	gagttcggga	cgTTTTgttat	tggtgagggt	attatgttgg	acgatagtgt	1320
tatggacgag	atagagttgg	gtggggaaaa	tatgttttgt	gatgataggg	gggtgacgtt	1380
tatagagttg	aagaaggggg	agtttttggt	ggataggagg	gcggatgtaa	ggtttttggc	1440
ggtttagttg	ttgttaggaat	agttttgggt	tatgtcgtcg	taggggttga	tgagtttatt	1500
gaattgcggg	tcgaagtatt	ttttgtatag	ttcggtttgt	tggttgcggt	TTTTTTTTTT	1560
ttatttggtt	cgacgatttt	tgaattaaat	ttgggggcgg	ttggggtaag	ggagtaaata	1620
gatgttatag	tgtagattat	taaaattttt	atcggagggt	aatttttcgtt	TTTTttcgat	1680
atatacgtaa	gcgtattttt	atattttggt	ttcggttttt	tgtatcggtt	tgtatattaa	1740
gatattaggg	ttagtttttt	gttattggtt	cgggttttta	ttaagcgtag	gagatttggt	1800
ttgttttggt	ttgcgagttg	ggattcggag	ttacgttata	aatttttagtc	gaacgtatgg	1860
agatttgcgg	acggtttgat	tatttagtta	ggcgtttttt	taggttttaa	aatatttaat	1920
gtaaaataaa	cgcggggtag	taggtttttt	taattttttt	cgggggtattt	tgtaaatttg	1980
tttttatttt	aaagttatag	atttacggat	gaggagaagg	gggtggaagg	gtattagagg	2040
atcgtttttt	tttttacgta	attttttttt	tttttttttg	atttttattg	tcgtttttta	2100
ttttttggtt	cgtgtttttt	taatagggat	taggtcggtt	atattttttt	tcgttttagta	2160
aaataattaa	ataaagagta	aaagattatt	ttttcggttag	ttcggttaatt	ttaggagttt	2220
ggtatatata	atttcgggaa	ttcggaagg	gtagtttttg	agattttttt	ttttttcggt	2280
ttgttttttt	tttattttaa	gtttattata	ggtttggtcg	cgcgttaggt	ttagtcgggt	2340
cgttttggtt	tgtaggcggg	tatttaggtc	ggtcgggttt	tattcgtgtt	cggtggttta	2400
gtcgtaatat	cgattttta	ttatatcggg	tttttttggt	gttttagacg	gcggtttttg	2460
tgtattggag	agaggttttg	tttgagatat	tcgagttgat	attagtgatg	ttttatatta	2520
tatattttcg	tcgggttttag	tcgtgttaatt	cgtttttttt	tttttttttt	ttatttttga	2580
tttttttttt	attttttttt	ttttttgtat	tcgattgtta	taaaaagtag	gttttatatt	2640
tatttggttc	gataagtagt	cgtttttggaa	ggagaggtag	ttgtaaggag	agtttagcgt	2700
cgcggttata	aagtattagg	gtggaggtgc	ggaatagcgg	gcgggggtggg	agggcgtttt	2760
cgaaggattt	tagaaaattt	atagattttg	tttttaatta	tttggtattt	ttatttttagg	2820
ttattttaaat	tttggttttagg	cgagaagagt	acgtgagagg	ttcggtttttt	tgatgtgtaa	2880
gagagttaat	gaaagattga	ttttgttttaa	aattacgtcg	tttaggattt	agtttttggtt	2940
ttggatagtt	aaattaaaat	tatttttaaat	tttttttcgg	ttttttattt	attagtatag	3000
ttttatgttt	tgtataaatg	ttatttagag	agtgtttttt	ttttttttga	tttgggagag	3060
tatttttggtt	tttatttttt	ttatcgttgt	tttttttttt	ttgtttgttt	tgtttttaate	3120
gggggttttt	ttttttttat	ttagagttat	taattttttt	tttttaatat	taaagttttt	3180
ggatgtcgtt	tgatttggtt	gattttgttt	tttgtttttt	gaattttta	aaatttgga	3240
tttttttatcg	attagtataa	attaggacgt	tgttattggg	ttattttatt	gagttttatt	3300
ttgttaatttt	ataaagtata	gattttgttat	aaagttaagg	taagtttttt	ttataaaatt	3360
atgattataa	tttagaagag	ggggtgtgag	ttttaatttt	tagagtttaa	tttttgagag	3420
aagataaata	aattaagtag	aaaagttttt	tttttttttt	tttttttttt	tttaagagga	3480
ttagtagttg	tgtattaaaa	ttttgttttc	ggagattata	aaattaggaa	atagggtgtg	3540
tgggagagat	ttgaatgggc	gaaataatcg	taaagaagg	gtaagaagcg	cgagtttagg	3600
agggaaaaag	ttgggttagg	gtcgggataa	aggtttttta	gggagggtta	attttttcgt	3660
gtttttggcg	ggtttttttt	gttaaagggt	tatagggttg	agtttggtcg	cggtttttgg	3720
tttggtaggg	attttattag	ttttgttttg	gtaattgtaa	gttaggaata	taatgttttg	3780
tgtaggggat	tgtttatgta	gttttagttcg	tgagatcgcg	ggatggcggg	gtagtgagtc	3840
gggtgtcgtt	tgggagtttg	agttaggggc	gtagttttgt	cggtttcgga	gaggggaattg	3900
taatttcgta	attaggtcgt	cgcgagggtt	tttggttttg	ttaaagttgcg	ttttatcggc	3960
gttttttttag	gcggcgttgt	tttttatatt	tttttttggt	ttatttggtt	tgtattttta	4020
taatattttt	tttttatatt	tttttagattt	cgtgttggtt	tttattcgga	ttcgggtttt	4080
cgtaagggtg	gtttatatag	cgattttttc	gcgtgtggat	atgttcgggt	agcgggtttt	4140
ttggaaagtg	gttttttaggt	tttgaggttg	ttgggttggt	aagtgaagtc	gttggtcgtt	4200
ttgtcgtttt	tttttagacg	ggtttttcggc	gtttacgttt	ttattttttt	tttggttggtt	4260
tttatttttt	tttgaaaacg	aaataatata	attttttcgt	tagtatgttt	atttgtaacg	4320
cggacgttaa	ttggatcggc	ggtagaagtc	gtggaagagt	tgggttggtt	ggcgtcggag	4380
gagggtgcgc	gcggcgggtt	cgggtcgcga	ggagcgttgc	gtttgtgggg	tgtgtaggcg	4440

taagtgtggg	tggttcgcggt	ttatTTTTTT	TTTTTTTTTT	gcgtcgtacg	TTTTatttat	4500
atgtttatTT	tattgtagcg	gtatatTTTt	TTTTatagtt	tgtgtTTTTt	agtatatTTt	4560
tatatTTTTg	cgtagatata	ttaaatTTTT	tgggacgcgt	atacgcgcg	ggtttataga	4620
TTTTTTTTTT	tttcgtagaa	agtttagatt	tttatgcggt	ttgggaaggt	taggaaaaga	4680
tgtgggggatt	cgggtgggta	tcgaagttcg	tcggTTTTTT	tttaaaaaaa	aaaaaaaaaT	4740
gtTTTTTTcgc	gaaggggtatt	tttgagtgg	tttaggtaat	TTTTtaacga	gtggagTTTT	4800
tcgggagttg	aaagtcgaga	ggaaaatagg	gatagaggtc	ggcggTTTTT	gaaggTTTTc	4860
gaattaagat	gttgggattt	ttgtgattt	ggaaatagaa	gggagggttag	ggtacgaata	4920
gagagggcg	tagaattggt	cgcgtTTTTt	gcgtTTTTtag	agtcgggtcg	gtcgagggag	4980
aattaaaggg	atgcggggta	gttaaaaatt	cggTTTTtcg	aagTTTTtgc	gggaggttag	5040
cgaacgatta	TTTTtattac	gtTTTTTTTT	ggagggggtg	atTTTTTTtg	ggcgagaggg	5100
agcgggtggc	gtagagtagt	tgagcgggaa	tgTTTTgtag	gcggcgcggc	gttttatttg	5160
cggTTTTtcg	gttggagggtg	tcggagatgg	tgtgtatTTT	tagtttgtgt	ttggaggagt	5220
ttagcgatcg	gggttgatcg	ggagttagaa	tcgaagtTat	ggttaacggt	tggggatgg	5280
gataggaaga	tgaggagacg	gtcgatagtt	tggTTTTcgt	tggtcgggtg	tttaagtga	5340
gcgggTTTTT	tatgtagttt	atggacgagg	gagcgcgcac	TTTTattagt	TTTTgggtat	5400
tgTTTTcgtc	agTTTTcgt	gtcgtcgttg	ttcgtttcgg	gtcgcgtTTT	aggcgcggag	5460
TTTTTTcgtt	gcggggagag	ttaggggacg	taatTTTcgt	cgagTTTTt	agttaagttg	5520
TTTTcgtTTT	tttcggaagg	tttaagcgaa	aaagtTcgga	gacggaaagt	tagcgggtaa	5580
acgaagatat	gggatgtggg	tagaagggt	ttatttagag	cgtTTTTtag	gagtaggTTT	5640
ttaagtTTTt	aagcgaaata	agagtgggt	aagatTTTTT	TTTTTTTTTT	TTTTTTTTTT	5700
taagaattTT	tttaataagg	aaagttaacg	tcgatcgcgt	tttgTtcgtt	TTTTTTTTt	5760
gcggtagttt	tgatagagaa	gtgttaagag	tgatagggat	aggtaggtga	tattagattt	5820
tttgcggcgg	tagtagtcgt	tgtagttacg	acgcggTTTT	ttgagcgtat	TTTTcgtaac	5880
gcgtatacgt	atTTTTTtcg	ggcggTcgaa	taggagtcgg	gttttgtcgt	agtttagttt	5940
taggtattt	ggcgagcgac	ggattagatt	tgcggtttcg	cgtTTTTTTg	ttggtttaat	6000
attttaaaaT	tagaggcggg	TTTTTTggtg	tcgagacgtt	atttcgtcgc	ggTTTTTTTT	6060
agTTTTTTTc	gttttcgttt	TTTTTTtagat	TTTTTTTcgg	gtgcgattga	cgtggTTTc	6120
tattaattag	gacgtTtcga	gtcgcgggtg	agggattggt	ttgTTTgtat	ttattagtag	6180
tgcggggtcg	ggttattggt	tcgtcgtgcg	tattgggttt	atataggtaa	gttttcggga	6240
atttagtttt	tgTTtagttt	aaggcgattc	ggTTTTtagt	acgaatttaa	aggTgaagag	6300
atgaggTtag	gagtcgaagg	tttgggagaa	gagagtggaa	tggTtaagaa	gagaaaggta	6360
taaggattaa	taagatatTT	attTTTTgtg	TTTTattata	tttattTTTt	attTTTTatt	6420
ttatataaaa	aggagatacg	ttatttaaaa	ttagaaaatt	tgaaaaatag	taataaatta	6480
TTTTTTcgat	tttaaatTTT	ttaaatagtt	tgTtaagtga	atgtTgcgtt	aattTgaaga	6540
agTTTTtaatt	gtaaagaaga	tagagtTTTg	aaaaggtagg	ttataaaatt	agaaatcgag	6600
aagTaaatgg	attcgTtaaa	agaaaattat	tttgattTTt	aacgaataat	tgTTTggTgg	6660
tttattTTTg	atttatataa	gaataaaaag	tcgtTTTtaga	ttacgtTTTT	tgtgatgttt	6720
attagttTTT	agatagaaaa	tatataatag	aagagaaatt	ttaatTTtag	gtTTTTtaaa	6780
tgttgaaagt	ttatttatTT	tattttaacgt	tgattaaagt	atatattTTt	gattTTTTta	6840
attTTTTgt	tattgtatta	agTtcgtTTT	aattcgcgag	agTtacgtTT	taaattcgat	6900
TTTTTTgttt	attttattat	taatttagatt	taaatttata	aagTTTgtag	aattaataat	6960
tttgagttaa	ttatatatga	aatatgtTTT	aatgaattTT	tatataatta	agaatgtTgt	7020
taaataatta	attTTaagga	taattTTTta	tagttattTT	TTTTTTTTtag	tgagtTTaag	7080
gttgTTTTga	gttattaaag	tttaagtagg	tagaaggggt	gtgtgtgagt	taagggcgaa	7140
aagTTtagaa	ttgcgtTTta	ttagTaaaag	taaaattTTt	tttatataaa	ataaaaaaaa	7200
ttattTTTg	agatatTaat	TTTTtatagt	attgtTTTt	agTaaattTt	attTTtaag	7260
aaattaaaga	aagaaattTt	aatatatTTt	aaataattTT	tgaaagTTTt	tttgTTTTTT	7320
agtataggTt	agTtgagag	gataaattTt	TTTTTTTTTg	gtTTTTgtat	gggcgattgt	7380
tttattatgg	agTtagtgTt	attattTTTg	aatgtgtatt	tgTTTgatat	tatagtTaat	7440
gattTgtaat	gttagtatga	agtattTTTt	aaatatTTTt	TTTTTgtTTT	tgTTTataag	7500
attgggaaat	ttattcgatg	tggaaataaag	tggatgaagt	agattataaa	tatatTTgt	7560
atttatgtgt	TTTTTTTTTg	TTTTgattat	TTTTaaattT	tattTgtaat	TTTTTTTTt	7620
TTTaaattTg	tagTTTaaag	acgtatatga	gaattgtTTT	ttagTTTTTT	TTTattagta	7680
ttattTTtatt	TTaagaataa	TTtagTTgt	agggaggaat	TTTTTTatag	taagTTTTta	7740
attagTattT	TTgtTTTTta	TTTTTTattT	tattTTattT	tattTTatat	atatagatat	7800
TTgtTTtagag	Taaaatatat	TTTTatgtga	taggtTTgt	ttagTTgagg	TTtatatat	7860
tagTTtatatt	aggTTTgt	attTTattat	Taaattatat	atattatat	agtagTTTgt	7920
TggTaaagaa	ggTtaaatta	atttatattT	TgtTTattat	TTggTgtTTt	aatgacgtat	7980
TTtattTtcg	agattTggcg	gagaattTTT	TTTTtagatt	TTatagcgtt	TTattgaaga	8040
TaatgtTTTT	atattTgtag	TggTTTTTaa	TTtgataaga	TTTtaattTg	TTtaagTTTt	8100
TTaaataagg	gtTTTaaatg	TTTTtagtcg	TTTTTTtatt	gaattTTTTT	TaatTTTTTT	8160
aagattataa	agtatatgtg	TaaagTaaat	attTTTTTTTT	attgtattgt	tagtcgatga	8220



tttataatta	agttaataag	aatttagttt	ttttttgttg	aatgtgttta	ttaattatat	8280
tttagttttt	ttttttaaat	tttagaatag	ttgtgggttt	tataatatta	tgttttttta	8340
agtttttatt	tatgaaggga	ttttattata	ttaaagaatg	aaaaaaattt	ttattgtagt	8400
tagtatatat	agttttttat	tttttgtttt	ttaagattta	aatttttagag	ttgtaaatat	8460
ttttggaagt	ttgggtgtta	atgtttttatt	ttagaaagtc	gagaagtttt	atagagttat	8520
atagattttt	aaattttatt	tttataaatt	tatagaattt	tgataaaaagt	tttgggtggtt	8580
ttatttttat	gatggaattt	ttattacgat	aaatatatat	gtatgaagga	ttttaattag	8640
tttttaaagt	ggttgaaaaa	tttaagggta	cgtgattggt	ttttatagtg	ttaacgtgtg	8700
cgagatgttg	gaagtattgg	ggattagtag	tagtttagat	gtttaaaaag	ataaggtggt	8760
ttaatttgtg	tggattttatt	gaagttaagt	ggtgaataaa	gataattatt	tagataattt	8820
agattaaagt	aaaagtaaaa	ttatatattt	ttgtatatat	atattttatat	ttattttata	8880
ttatagatat	atatacgtat	atatatattg	gttttgtaaa	taattgattt	aaagtgagga	8940
ttttttttgt	attttttttag	taggagtttt	aatatttttt	taatttttta	attattttat	9000
a						9001

<210> 3

<211> 9001

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 3

tgtaaagtga	ttaggagatt	aggagaatgt	tgaaattttt	gttggaaaaa	tgtaaagaaa	60
attttttatt	tgagtttagt	gtttatagag	ttagtgtgtg	tgtgcgtgtg	tgtgtttgta	120
atataaaaatg	gatgtgaata	tatatatata	aatagatatg	gttttgtttt	tatttttaatt	180
tgaattattt	agataattgt	ttttatttat	tatttgattt	taatgggttt	atataaatta	240
ggatatttta	tttttttagg	tatttagggt	gttggtgatt	tttagtggtt	ttaatatttc	300
gtatacgttg	gtattatgag	gagtagttac	gtgttttttg	gttttttaaat	tattttggag	360
gttgattgag	gtttttttata	tatgtatat	tgtcgtgatg	aaagttttat	cggtagagtg	420
gagttattag	agttttttatt	aaaattttgt	gggttttatga	gagatgggtt	tagaaattta	480
tatgggtttt	tgggggtttt	cgggtttttta	aaataaggta	ttaatattta	agtttttaaa	540
aatatttgta	gtttttgggt	ttgaattttg	aaaaataagg	agtgaagggt	tgtgtatat	600
aattatagtg	gagatttttt	ttatttttta	atgtgatgga	gtttttttat	gaaatgaagt	660
tttaaggggt	atgggtattgt	ggggattata	gttattttga	ggtttaaaag	aagaaattgg	720
aatatgatta	gtaaatatat	ttagtagaaa	agagttggat	ttttattgat	ttagttatag	780
gttatcgggt	ggtagtgtaa	tgggaggaaa	tattttattt	atatatatat	tttatgattt	840
tgggggaatt	agaggaaatt	taataagaaa	acggttagaa	atatttaaaa	tttttattta	900
aaagatttaa	gtaaattaga	gtttttattag	attaaaaatt	attataaatg	taagagtatt	960
gttttttagtg	aaacgttggt	gggttttgaga	aggagatttt	tcgttaaatt	ttcgggataa	1020
aatgcgttat	ttaagtatta	gataatgagt	agaatgtaaa	ttaatttaaat	tttttttatt	1080
aatagggttg	tagtgtaaat	tgtataattt	agtataaga	ttgtaggatt	taatataagt	1140
ggatgtatga	gttttagtta	atgtagattt	gttatatgag	gatgtgtttt	attttgagta	1200
ggtgtttgta	tgtgtggaat	ggggtaaagt	ggaataaaaag	gttaaaaagta	gaaatgttga	1260
tttaaagttt	attatgaaga	aatttttttt	ttgtagttaa	attattttta	aagtgggatg	1320
atattggtga	agaaagattg	aaaaataatt	tttatgtgcg	tttttggatt	gtaagtttaa	1380
aatggggagg	agttgtagat	agggtttggg	ggtggttagg	gttaaaggaga	gatataataag	1440
ttgtaaatat	atttgtagtt	tgttttattt	attttgtttt	atatcgaata	agttttttta	1500
ttttgtgaat	aaggataagg	agggagtgtt	ttaaagatat	tttatgttgg	tattgtaaat	1560
tattgattgt	aatgttaaat	aaatatatat	ttagagatga	taatattaat	tttatagtaa	1620
aataatcgtt	tatgtagaaa	tttagaggag	attagtttgt	tttttttagt	tgatttatgt	1680
tgggggataa	aaggattttt	aaaaattatt	ttgaatatgt	ttggattttt	tttttttaatt	1740
tttttggaag	ttaaatttgt	ttggaaatag	tgttataaag	agttgatggt	tttaaagggtg	1800
attttttttg	ttttatataa	ataagggttt	gtttttgtta	gttgagcgta	gttttagggt	1860
tttcgttttt	agtttatata	tatttttttt	gtttgttttg	attttaaatg	tttaagatag	1920
ttttgagttt	attgggaaaa	gaaaatgatt	gttaaaaaatt	atttttgaaa	ttgggttattt	1980
ggtaaatatt	ttaattgtat	ggaaatttat	taaggatatat	tttatatata	attagtttaa	2040
ggttggtgat	tttatagggt	ttatggattt	aaatttgatt	gataataaag	ttaaataagag	2100
agtegaattt	aaagecgtgt	tttttcgggt	taggacgagt	ttaatatagt	gtataaggaa	2160
tttgaaagat	ttaggatatg	tgttttaatt	aacgttaagt	agaatggata	agtttttagt	2220
attttgaaaa	cgttgggtta	gggttttttt	tttattgtgt	gttttttgtt	tggggattaa	2280

taagtattat	agagaacgtg	at ttgagggcg	at ttttttatt	tttgtataaaa	tttagagtga	2340
attattaaat	agttgttcgt	ttaaagttaa	ggtaat tttt	ttttgacggg	tttatttggt	2400
tttcgatttt	taattttatta	gtttgttttt	ttaggggttt	gtttttttttg	taattaaagt	2460
tttttttagat	tagcgtagta	tttattttgat	aggttgtttg	gaaaatttaa	gacgagagag	2520
gtgatttggt	gttgtttttt	aaat tttttta	gttttaagta	acgtgttttt	tttttatatg	2580
gggtggggga	ttggaaatgg	atgtagtgag	atataaagag	tgggtgtttt	gttgattttt	2640
gtattttttt	ttttttgatt	at tttat ttt	tttttttttaa	gttttcgatt	tttagt tttta	2700
ttttttttatt	tttgggttcg	tattaaaagt	cggatcgttt	tgggttgggt	aggagt t gaa	2760
ttttcgggag	tttgtttgtg	tagattttagt	gcgtacggcg	aggtagtagt	tcggtttcgt	2820
attgttgata	ggtgtaggta	ggatag tttt	tttatcgcgg	ttcggggcgt	tttgattggg	2880
gcggagttac	gttagtcgta	ttcggagaag	ggtttgggag	gaggcggagg	cggagagggg	2940
tggggagggg	cgcggcggag	tgacgttttcg	gtattaggaa	gttcgttttt	ggttttaaga	3000
tgttaggtta	ataggggaagc	gcggagtcgt	agatttggtt	cgtcgttcgt	ttgggtgttt	3060
ggagttgagt	tgcggttaagg	ttcggttttt	gttcgatcgt	tcgaggggtg	tgcggtg tgcg	3120
cgttgccggag	ggtgcgttta	gagggtcgcg	tcgtggttgt	agcggttgtt	gtcgtcgtag	3180
gggatttaaat	attattttatt	tgtttttgtt	at tttttgata	ttttttttgtt	agggttg tgcg	3240
cgtggggggg	gggcgggtag	agcgcgg tgcg	gcgttagttt	tttttattgg	agggg ttttt	3300
gggggagggg	gggagagaag	aaggggg ttt	ttgtttattt	ttgtttcgtt	ttggagt tttg	3360
gaagt tttgtt	ttttaaagac	gtttttgagt	gtgttttttt	gtttatatatt	tatgt ttttcg	3420
tttgttcgtt	gattttttcgt	tttcggattt	tttcgttttg	gttttttcgga	ggagacgggg	3480
gtagtttggt	ttgagaattc	ggcgggggtt	gcgtttttttg	gttttttttcg	tagcggggaa	3540
at ttcgcgtt	tagagcgcga	ttcggagcgg	gtagcggcgg	ttacgggggt	tcggcggggg	3600
agtagttaag	gattagtaga	gcgtcgcgtt	ttttcgttta	tgaattgtat	gaaagg ttcg	3660
ttttattttgg	agtatcgagt	agcggggatt	aagttgtcgg	tcgttttttt	at tttttttgt	3720
tattattttt	agtcgttagt	tatggttttcg	gttttggttt	tcggtttagt	tcggtcgttg	3780
gatttttttta	agtatagggt	ggaggtgtat	attatttttcg	atattttttag	ttcggagggtc	3840
gtaggttaagg	cgtcgcgtcg	ttttgtagat	at ttttcgttt	agttgtttttg	cgttattcgt	3900
ttttttttcgt	tttaagggaag	ttagtttttt	cgggggggagg	cgtgggtggga	gtgggtcgttc	3960
gttttggtttt	tcgtagaatt	ttcgggagtc	ggaatttttg	ttattttcgt	ttttttttagt	4020
ttttttttcga	tcggtttcgtt	ttttggggcg	ttaagggcgc	gagtaatttt	gtcgtttttt	4080
ttattcgtat	tttggtttttt	tttttggttt	ttgggttata	aaaatttttag	tatttttgatt	4140
cgaggatttt	tagaggtcgt	cgattttttgt	ttttgttttt	ttttcggttt	ttagt ttttcg	4200
aggagt tttta	ttcgttagga	aattgttttg	aattattttag	aaatgttttt	cgcgaagagg	4260
tattttttttt	tttttttttg	gaaagggtcgc	gcgaattttcg	gtgttttaatc	gaattttttat	4320
at ttttttttt	agtttttttta	aatcgtatgg	aaattttgagt	tttttgcgag	ggggaggggg	4380
gtttgtaaat	tacgcgcgtg	tgccgcgtttt	aggagattttg	gtgtgttttgc	gtagaggtgt	4440
ataaatatat	ttgaaagtat	aggttataaaa	agtgaatgtg	tcgttgtagt	gagataaata	4500
tgtaaaataaa	acgtgcggcg	ttggggggagg	ggaggaaatg	gggcgcggat	at tttatat t	4560
gcgttttgat	at ttttatagg	cgtagcgtttt	ttcgcggttc	ggagtcgtcg	cgcgtat tttt	4620
ttttcggcgt	taggtagttt	agtttttttta	cggttttttgt	cgtcggttta	gttggcgttc	4680
gcgttgtagg	tgggtatgtt	gacgggaaag	tgtgtgtgtt	tcgttttttag	agaaagataa	4740
aagttagtag	gggaagaatg	aggacgtggg	cgtcagggat	tcgttttaaga	agaagcggta	4800
aaggcggtag	cggattttatt	ttatttagtta	gtagtttttag	gagttggagg	ttattttttta	4860
gaggaatcgt	tattcggata	tgtttatacgc	cgaagaaatc	gttgtgttga	ttaat ttttac	4920
ggaagt ttcga	gttcgggtag	gagttagtag	ggagt tttggg	agggatgggg	ggaggatgtt	4980
gtggagggtat	aggttaagta	gattaggaga	gaatgtggaa	ggtagcgtcg	tttgggaggg	5040
cgtcgggtggg	gcgtagtttt	gtaaaggtag	aaggttttcgc	ggcgggtttgg	ttgcgagatt	5100
atagttttttt	tttcgagggtc	gataggattg	tcgtttttggt	ttaggttttt	agagcggtat	5160
cggttttattg	tttcgtttatt	tcgcgatttt	acgagttggg	ttgtatgggt	aatttttttgt	5220
ataggatatt	gtgttttttg	tttgtagttg	ttagagtaga	gttaataaaa	ttttttattag	5280
gttaagagtc	gcgaatagggt	tttaattttgt	gagtttttaa	taaggaaaat	tcgttagaga	5340
tacggaagag	ttggtttttt	ttgggaaatt	tttgttttcgg	ttttgggttta	gtttttttttt	5400
ttttgggttc	gcgttttttta	tattttttttt	acggttggtt	cggttatttta	ggttttttttt	5460
atataatttta	tttttttagtt	ttgtgatttt	cgggagtaaaa	gtttttaatat	ataattatta	5520
gttttttttag	aaggagaaag	aaaaaaagaa	gaaagatttt	tttggtttggt	ttattttattt	5580
tttttttagga	gttgaatttt	ggaaatttgaa	at tttatat t	tttttttttaa	attataatta	5640
tagtttttgta	aaaagggttt	at ttttaattt	tgtagtaaat	ttgtattttta	tggatttggt	5700
aaaatgagtt	taaataaata	at tttaatagt	aacgtttttgg	tttatgtttg	tcggtggaag	5760
at ttttaaat	tgttaggatt	ttggaagtag	aaaatagaat	taagtaaat	aagcggatt	5820
tagaggtttt	gttgtttaaaa	aaaaaaaatt	aagtgttttg	ggtagaaaaa	ataaagtttt	5880
cggttagagt	agagtaaaata	aaaagaagaa	aataacgata	aaaagaataa	agattaaaat	5940
gttttttttaa	at tagagggg	atgaagatat	tttttggtg	gtattttgtgt	aaggtatgag	6000
gttatgttgg	tggataaaaag	gtcgggaaga	agttgaaaat	ggtttttagtt	taattgttta	6060



gagttagagt	tgggttttgg	gcggcgtggt	tttgagtaag	gttagttttt	tattagtttt	6120
tttgtatatt	aagggaacgg	gtttttttacg	tattttttttc	gtttgagtaa	agtttagatg	6180
gttttagggt	gaaatggtaa	gtaattaaag	atagagttta	tgggtttttt	gggatttttc	6240
gaaaacgttt	ttttatttcg	ttcgttattt	cgtagtttta	ttttagtgtt	ttgtagtcgc	6300
ggcgttgggt	tttttttgta	gttggttttt	tttttagggc	ggttgtttgt	cgagttaagt	6360
gggagtgagg	cgtgtttttt	atagtagtcg	ggtgtaaaga	ggaaggggga	taaaaaggaa	6420
attaagaatg	aaaggaaaaa	gagaaaaagc	ggattatacg	gttgggttcg	gcggagatgt	6480
gtaatgtgaa	atattattgg	tgtagtttcg	gatatttttag	gttaggtttt	tttttaatat	6540
ataaaagtcg	tcgtttgggg	cgatagggag	gttcgatgtg	gattgggatc	ggggttgcgg	6600
ttgggttatc	ggatacgggt	ggaagtcggt	cggtttgggt	ggtcgtttgt	aaagttaaag	6660
gattcgggtg	ggtttggcgc	gcggataggt	ttgtgggtgg	tttagggtaa	agaagaggta	6720
gagcgaaaga	agggggaatt	tttaaaatta	tttttttcgg	gttttcggag	tttaatatgt	6780
taagtttttg	gagttaacga	gttgacgaag	aggtgggttt	ttgtttttta	tttgggtgtt	6840
ttgttagggc	agaaagagt	ttggcgggtt	agttttttgt	aaggagtagc	gtattagggg	6900
gtgggggacg	atagtggagg	ttagggaagg	aaggaggaga	ttgcgtggga	gaaagagcga	6960
tttttttagt	tttttttagt	tttttttttt	tattcgtggg	tttgtgtttt	tggaatggaa	7020
gtaagtttgt	aaggtgtttc	gggaagggtt	ggaaaagttt	gttgttttcg	gtttgtttta	7080
tattaagtgt	ttttggattt	ggagaaacgt	ttgggttgagt	gattaaatcg	ttcgtagggt	7140
tttatgcgtt	cgggttgaggt	ttgtggcgta	gttttcgaggt	ttagttcgta	ggttagagta	7200
gattaggttt	tttgcgtttg	gtggagattc	gggttagtaa	ttgaaagtgt	gttttggtat	7260
tttgggtgtg	agggcgggtg	agtgaagcga	ggttaggggt	tgtgagtgcg	ttagcgtgtg	7320
tgtcggggga	aggcgggggt	tggttttcga	tggaagtttt	agtaatttgt	attgtggtat	7380
ttgtttgttt	ttttgtttta	atcgtttttt	ggtttggttt	aagaatcgtc	gggttaaagt	7440
gagaaagagg	gagcgttaatt	agtaggtcga	gttatgtaag	aatggtttcg	ggtcgtagtt	7500
taatgggttt	atgtagtttt	acgacgatat	gtatttaggt	tattttttata	ataattgggt	7560
cgttaagggt	tttatattcg	ttttttttatt	tattaagagt	tttttttttt	ttaatttttat	7620
gaacgttaat	tttttggtat	tatagagtat	gttttttttta	tttaatttta	tttcgttttat	7680
gagtatgtcg	tttagtatgg	tggttttagt	agtgataggc	gttttcgggt	ttagttttaa	7740
tagtttgaat	aatttgaata	atttgagttag	ttcgtcgttg	aatttcgcgg	tgtcgacgtt	7800
tgtttgtttt	tacgcgtcgt	cgattttttc	gtatgtttat	agggatacgt	gtaattcgag	7860
tttggttagt	ttgagattga	aagtaaagta	gtatttttagt	ttcgggttacg	ttagcgtgta	7920
gaattcgggt	tttaatttga	gtgtttgtta	gtatgtagtg	gatcgggttcg	tgtgagtcgt	7980
atttatagcg	tcgggatttt	aggattttgt	cggatgggggt	aatttcgttt	ttgaaagatt	8040
gggaatttat	ttagaagggtc	gtgggtatta	aagaaaggga	gagaaagaga	agttatatag	8100
agaaaaggaa	attattgaat	taaagagaga	gttttttttga	ttttaaaggg	atgttttttag	8160
tgtttgatat	tttttattat	aagtattttt	aatagttgta	aggatatata	tataaataaa	8220
tgtttgattg	gatatgatat	tttaaatatta	ttataagttt	gttattttttt	aagtttagta	8280
ttgttaatat	ttaaatgatt	gaaaggatgt	atatatatcg	aaatgtttaa	ttaatttttat	8340
aaaagtagtt	gttagtaata	ttataatagt	gttttttaaag	gttaggtttt	aaaataaagt	8400
atgttatata	gaagcgatta	ggattttttcg	tttgcgagta	agggagtgtg	tatattaaat	8460
gttatattgt	atgttttttaa	tatattatta	ttattataaa	aaatgtgtga	atattagttt	8520
tagaatagtt	tttttggtgg	atgtaatgat	gttttttgaaa	ttgttatgta	taattttattt	8580
tgtgtataat	atttcgtata	atattattgt	tttatttttt	agtaaatatg	aaataaatgt	8640
gttttatttt	atgggagtaa	aatatatatt	atataaattg	gttttggttt	ttttttttttt	8700
tttttgttat	taatttggtt	aggatatatt	agttattgtt	ttttaaataa	attagttttt	8760
tttgtttgtt	tagtttaaata	tataaggtag	tagttttttat	ttaaatttgg	tagaaataaa	8820
tgatagttat	ttattagaaa	ttaaaaagaa	aaaaaaaggt	attttcgggg	gggaaaaggg	8880
ttataaaaatt	taattttgtt	tttttaattt	ttttttggtt	taaatttaga	ggattttatt	8940
atggttagta	aataatatga	aaaagaaaaa	agaagaaaga	aatttagtaa	gtttattagt	9000
t						9001

<210> 4

<211> 9001

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 4

agttgatgga	tttgttaaatt	ttttttttttt	ttttttttttt	tttatattat	ttgttagtta	60
taatggaatt	tttttaggtt	aagttaaaga	aaaattggag	agataaaaatt	agatttttga	120

gtttttttttt	tttttgggaa	tgttttttttt	tttttttttta	gtttttgatg	aatgggttatt	180
attttattttt	attaaatttta	aataaggatt	gttgtttttgt	atgttttaatt	aggtaggtag	240
aggggaattgg	tttgtttagg	aagtagtgat	tgagatgttt	tggttaagtt	agtgatagag	300
gaggggagaa	agaatttaga	ttaatttgta	tgtagtatat	tttatttttta	tgaaataaaa	360
tatatttggt	ttatatttgt	tgaaaagtaa	aataataata	ttgtatgaaa	tgttatatat	420
agggtagggt	gtatatagta	gttttagaaa	tattattgta	tttattagag	aaattattttt	480
aaaattgata	tttatatatt	ttttataata	ataataatat	gttagaaata	tatagtgtgg	540
tatttagtat	atataattttt	ttgtttgtaa	gtgaaaaatt	tttaattgttt	ttgtataata	600
tgttttatttt	taaagttttaa	tttttaaaaa	tattgttgtg	atattattaa	taattgttttt	660
tataaaaatta	atttgatatt	ttgatataata	tataatttttt	tagttatttta	aatgtttaata	720
atgttaaatt	taaaaaataa	taagtttata	gtaatgttaa	aatgttatat	ttagttaaata	780
atttgtttgt	gtatgtgttt	ttgttaattgt	tagaaatatt	tgtagtgaata	gatgttagat	840
attgaggata	tttttttgaa	attaaaggag	ttttttttttt	gatttagtgg	ttttttttttt	900
tttatatagt	ttttttttttt	ttttttttttt	ttagtgttta	tgattttttta	gtataattttt	960
tagttttttta	agggtggagt	tgtttttattt	ggtaagggttt	taggatttttg	gtgttgtggg	1020
tgtggttttat	atgggttggt	ttattgtata	ttggtaagta	tttaggttg	agggtgggtt	1080
ttgtatgttg	gtgtagttga	agttggagtg	ttgttttgtt	tttagtttta	gggtgggttag	1140
gtttgagtta	tatgtgtttt	tataaatata	tggaggagt	gggtgtgtgt	aaggatagg	1200
agggtgttgg	attgtggaat	ttagtgtatg	gttatttagg	ttgtttaagt	tatttaggtt	1260
gttgagattg	gagtttgga	tgtttgttat	tgttgagggt	attatgttgg	atgatagt	1320
tatggatgag	atagagttgg	gtggggaaaa	tatgttttgt	gatgatagg	ggttgatgtt	1380
tatagagttg	aagaagggga	agtttttggt	ggataggagg	gtggatgtaa	ggtttttggt	1440
ggtttagttg	ttgtaggaat	agtttggtga	tatgttgttg	tagggttgta	tgagtttatt	1500
gaattgtgg	ttgaagttat	ttttgtatag	tttggttgtt	tgggtgtgtt	ttttttttttt	1560
ttatttggtt	tgatgatttt	tgaattaaat	ttgggggtgg	ttggggtaag	ggagtaaata	1620
gatgttatag	tgtagattat	taaaattttt	attggagggt	aatttttgtt	tttttttgat	1680
atatatgtta	gtgtatttat	atattttggt	tttggtttat	tgtattgttt	tgtatatata	1740
gatattagg	ttagttttta	gttattgggt	tgggttttta	tttaagtgtg	gagatttggt	1800
ttgttttggt	ttgtgagttg	ggatttgagg	ttatgttata	aatttttagt	gaatgtatgg	1860
agatttggtg	atgggttgat	tatttagtta	gggtgttttt	taggttttaa	aatatttaata	1920
gtaaaataaa	tgtggggtag	taggtttttt	taattttttt	tgggggtatt	tgtaaatttg	1980
ttttttatttt	aaagttatag	atttatggat	gaggagaagg	gggtggaagg	gtattagagg	2040
attgttttttt	tttttatgta	attttttttt	ttttttttttg	attttttattg	ttgttttttta	2100
tttttttggt	tgtgtttttt	taatagggat	taggttggtta	atattttttt	ttgttttagta	2160
aaataatttaa	ataaagagta	aaagattatt	tttttggttag	tttgtttaatt	ttaggagt	2220
ggtatatata	attttgggaa	tttggaagg	gtagtttttg	agattttttt	tttttttggt	2280
ttgtttttttt	tttatttttaa	gtttattata	ggtttggttg	tgtgttaggt	ttagtgggt	2340
tgtttgggtt	tgtagggtgg	tatttaggtt	gggtgggttt	tatttgtgtt	tgggtgggtta	2400
gttgtaattt	tgattttta	ttatatggg	tttttttgtt	gttttagatg	gtgggttttg	2460
tgtattggag	agaggtttg	tttgagatat	ttgagttgat	attagtgatg	ttttatatta	2520
tatattttttg	ttgggttttag	ttgtgtaatt	tgtttttttt	ttttttttttt	ttattttttga	2580
ttttttttttt	attttttttt	ttttttgtat	ttgattgtta	taaaaagtat	gtttttatttt	2640
tatttggttt	gataagtagt	tgttttggaa	ggagaggtag	ttgtaaggag	agtttagtgt	2700
tgtgggtata	aagtattagg	gtggagttgt	ggaatagtgg	gtgggggtgg	agggtgtttt	2760
tgaaggattt	tagaaaattt	atagattttg	tttttaatta	tttgttattt	ttatttttagg	2820
ttattttaaat	tttgtttagg	tgagaagagt	atgtgagagg	tttgtttttt	tgatgtgtaa	2880
gagagttaat	gaaagattga	ttttgttttaa	aattatgttg	tttaggattt	agttttgggt	2940
ttggatagtt	aaattaaaat	tattttttaat	tttttttttg	ttttttattt	attagtatag	3000
ttttatgttt	tgtataaatg	ttatttagag	agtgttttta	tttttttttga	tttgggagag	3060
tatttttggt	tttatttttt	ttattgttgt	ttttttttttt	ttgtttgttt	tgttttaatt	3120
gggggtttta	tttttttttat	tttagagtatt	taattttttt	tttttaatag	taaagttttt	3180
ggatgttgtt	tgatttggtt	gattttgttt	tttggttttta	gaattttta	aaatttgga	3240
ttttttattg	attagtataa	attaggatgt	tgttattggg	ttattttattt	gagttttattt	3300
ttgttaattt	ataaagtata	gatttggtat	aaagttaagg	taagtttttt	ttataaaatt	3360
atgattataa	tttagaagag	ggggtgtgag	ttttaatttt	tagagttaaa	tttttgagag	3420
aagataaata	aattaagtag	aaaagttttt	ttttttttttt	ttttttttttt	tttaagagga	3480
ttagtagttg	tgtattaaaa	ttttgttttt	ggagattata	aaattaggaa	atagggtgtg	3540
tgggagagat	ttgaatgggt	gaaataattg	taaagaaggt	gtaagaagtg	tgagttagg	3600
agggaaaaag	ttgggttagg	gttgggataa	agggtttttta	gggagggtta	atttttttgt	3660
gttttttggtg	ggtttttttt	gttaaagggt	tatagggttg	agtttggttg	tgggtttttg	3720
tttggtaggg	attttatttag	ttttgttttg	gtaattgtaa	gttaggaata	taatgttttg	3780
tgtaggggat	tgtttatgta	gttttagttg	tgagattgtg	ggatgggtgg	gtagtgaagt	3840
gggtgtgttt	tgggagtttg	agttagggtg	gtagtttttgt	tgggttttga	gaggggaattg	3900



taatTTTTgta	attaggttgt	tgtgaggttt	tttgtttttt	taaagttgtg	ttttatttgg	3960
gttttttttag	gtggtgttgt	tttttatatt	tttttttgg	ttatttgggt	tgtattttta	4020
taatatTTTT	tttttatttt	tttttagatt	tgtgttgggt	ttatttggga	tttgggtttt	4080
tgtaaaggtg	gtttatatag	tgattttttt	gtgtgtggat	atgtttgggt	agtgggtttt	4140
ttggaaagtg	gttttttagt	tttggagttg	ttggttggta	aagttagttt	gttgttgttt	4200
ttgttgtttt	tttttagatg	gggttttgg	gtttatgttt	ttattttttt	tttgttgggt	4260
tttatttttt	tttgaaaatg	aaatatatat	atttttttgt	tagtatgttt	atttgtaatg	4320
tggatgttaa	ttggattgg	ggtagaagtt	gtggaagagt	tgggttgttt	gggttggag	4380
gaggtgtgt	gtggtgggtt	tgggttgtga	ggagtgttgt	gtttgtggg	tgtgtaggtg	4440
taagtgtggg	tgtttgtgtt	ttattttttt	ttttttttta	gtgttgtatg	ttttatttat	4500
atgtttattt	tattgtagt	gtatatattt	ttttatagtt	tgtgttttta	agtatattta	4560
tatatTTTTg	tgtagatata	ttaaattttt	tgggatgtgt	atatgtgtgt	ggtttataga	4620
tttttttttt	ttttgtagaa	agttagatt	tttatgtgt	ttgggaaggt	taggaaaaga	4680
tgtggggatt	tgggtgggta	ttgaagtttg	ttggtttttt	tttaaaaaaa	aaaaaaaaat	4740
gtttttttt	gaagggatt	tttgagtgg	tttaggtaat	tttttaata	gtggagtttt	4800
ttgggagttg	aaagttgaga	ggaaaatagg	gatagaggtt	gggtgttttt	gaaggttttt	4860
gaattaagat	gttgggattt	ttgtgattta	ggaaatagaa	gggaggttag	ggtagaata	4920
gagaggggtg	tagaattgtt	tgtgttttta	gtgttttagg	agttgggttg	gttgagggag	4980
aattaaaggg	atgtggggta	gttaaaaatt	tgggtttttg	aagttttgtg	gggagttagg	5040
tgaatgatta	tttttattat	gttttttttt	ggaggggttg	atttttttgg	ggtagagagg	5100
agtgggtgg	gtagagtagt	tgagtgggaa	tgtttgtagg	gtgggtgtgt	gttttatatt	5160
tgggtttttg	gttggaggtg	ttggagatgg	tgtgtatttt	tagtttgtgt	ttggaggagt	5220
ttagtgattg	gggttgattg	ggagttagaa	ttgaagttat	ggtaaatgg	tggggatgg	5280
gataggaaga	tgaggagatg	gttgatagtt	tgggtttttg	tgtttgggtg	tttaagtga	5340
gtgggttttt	tatgtagttt	atggatgagg	gagtgtgatg	ttttattagt	ttttggttat	5400
tgttttgttg	agtttttgt	gttgttgttg	tttgttttgg	gttgtgtttt	aggtgtggag	5460
tttttttgtt	gtggggagag	ttaggggatg	taatttttgt	tgagttttta	agttaagttg	5520
tttttgtttt	ttttggaagg	tttaagtga	aaagtttgga	gatggaaagt	tagtgggtaa	5580
atgaagatat	gggatgtggg	tagaagggt	ttatttagag	tgtttttagg	gagtaggttt	5640
ttaagtttta	aagtgaata	agagtgggt	aagatttttt	tttttttttt	tttttttttt	5700
taagaatttt	tttaataagg	aaagttaatg	ttgattgtgt	tttgtttgtt	tttttttttat	5760
gtggtagttt	tgatagagaa	gtgttaagag	tgatagggat	aggtaggtga	tatttagattt	5820
tttgtggtgg	tagtagttgt	tgtagtattg	atgtggtttt	ttgagtgtat	tttttgtaat	5880
gtgtatatgt	atattttttg	gggtgttgaa	taggagttgg	gttttgttgt	agtttagttt	5940
taggtatttt	ggtagagtgt	ggattagatt	tgtgggtttt	tgtttttttg	ttgggttaat	6000
atttttaaaat	tagaggtggg	tttttttgg	ttgagatgtt	attttgttgt	ggtttttttt	6060
agtttttttt	gttttttgtt	tttttttag	tttttttttgg	gtgtgattga	tgtgggtttt	6120
tattaattag	gatgttttga	gttgtgtgtg	agggattgtt	ttgtttgtat	ttattagtag	6180
tgtgggggtg	ggttattgtt	ttgttgtgtg	tattgggttt	atataggtaa	gtttttggga	6240
atttagtttt	tgttttagttt	aaggtgattt	ggtttttagt	atgaatttaa	agggtgaagag	6300
atgaggttag	gagttgaagg	tttgggagaa	gagagtggaa	tgggttaagaa	gagaaaggta	6360
taaggattaa	taagatatatt	attttttgtg	ttttattata	tttattttta	attttttatt	6420
ttatataaaa	aggagatatg	ttattttaaaa	ttagaaaatt	tgaaaaatag	taataaatta	6480
tttttttgat	tttaaatatt	ttaaatagtt	tgttaagtga	atgttgtgtt	aatttgaaga	6540
agtttttaatt	gtaaagaaga	tagagttttt	aaaaggtagg	ttataaaatt	agaaattgag	6600
aagtaaatgg	atttgttaaa	agaaaattat	tttgattttt	aatgaataat	tgtttgggtg	6660
tttatttttg	atttatataa	gaataaaaag	ttgttttaga	ttatgttttt	tgtgatgttt	6720
attagttttt	agatagaaaa	tatatataat	aagagaaatt	ttaatattagt	gttttttaaaa	6780
tgttgaaagt	ttatttatatt	tatttaaatgt	tgattaagat	atatatttta	gatttttttaa	6840
attttttgt	tattgtatta	agtttgtttt	aatttgagag	agttatgttt	taaatttgat	6900
tttttttgtt	attttattat	taatttagatt	taaatttata	aagttttag	aattaataat	6960
tttgagttaa	ttatatatga	aatatgtttt	aatgaatttt	tatatataat	agaatgttgt	7020
taaataatta	attttaagga	taatttttaa	tagttatttt	tttttttttag	tgagtttaag	7080
gttgttttga	gttattaaag	tttaagtagg	tagaaggggt	gtgtgtgagt	taaggggtga	7140
aagtttagaa	ttgtgtttta	ttagtaaaag	taaaattttt	tttatataaa	ataaaaaaaa	7200
ttattttttg	agatattaat	tttttatagt	attgtttttt	agtaaaattt	attttttaag	7260
aaattaaaga	aagaaattta	aatatatatt	aaataatttt	tgaaagtttt	tttgtttttt	7320
agtataggtt	agttggagag	gataaattaa	tttttttttgg	gtttttgtat	gggtgattgt	7380
tttattatgg	agttagtgtt	attatttttt	aatgtgtatt	tgtttgatat	tatagttaat	7440
gatttgtaat	gttagtatga	agtatttttt	aaatatatttt	tttttgtttt	tgtttataag	7500
attgggaaat	ttatttgatg	tggaaataaag	tggatgaagt	agattataaa	tatatattgta	7560
atttatgtgt	tttttttttt	ttttgattat	ttttaaatatt	tatttgtaat	tttttttttat	7620
tttaaatattg	tagtttaaaag	atgtatatga	gaattgtttt	ttagtttttt	tttattagta	7680

ttatttttatt	ttaagaataa	tttagttgta	agggaggaat	tttttttatag	taagtttttaa	7740
attagtatttt	ttgttttttaa	tttttttattt	tatttttattt	tatttttatat	atatagatat	7800
ttgttttagag	taaaatatat	ttttatgtga	taggtttgta	ttagttgagg	tttatatatt	7860
tagttatatt	aggttttgta	atttttattat	taaattatat	atatttatatt	agtagtttgt	7920
tggtaaagaa	ggtttaaatta	attttatattt	tgtttattat	ttgggtgttta	aatgatgtat	7980
tttatttttgg	agattttggtg	gagaatttttt	tttttagatt	ttatagtgtt	ttattgaaga	8040
taatgtttttt	atattttgtag	tgggttttttaa	tttgataaga	ttttaattttg	tttaagttttt	8100
ttaaataagg	gtttttaaatg	tttttagttg	ttttttttatt	gaatttttttt	taatttttttt	8160
aagattataa	agtatatgtg	taaagtaaatt	atttttttttt	attgtattgt	tagttgatga	8220
tttataatta	agttaataag	aatttagttt	tttttttggtg	aatgtgttta	tttaattatat	8280
tttagtttttt	tttttttaaat	tttagaatag	ttgtgggtttt	tataatatta	tgtttttttaa	8340
agtttttattt	tatgaaggga	ttttattata	ttaaagaatg	aaaaaaattt	ttattgtagt	8400
tagtatatat	agtttttttat	tttttggtttt	ttaagatttta	aatttttagag	ttgtaaatat	8460
ttttggaagt	ttgggtgtta	atgtttttatt	ttagaaagtt	gagaagtttt	atagagttat	8520
atagatttttt	aaattttattt	tttataaatt	tatagaattt	tgataaaaagt	tttggtgggtt	8580
ttatttttatt	gatggaattt	ttattatgat	aaatatatat	gtatgaagga	ttttaatttag	8640
tttttaaaagt	ggttgaaaaa	tttaagggtta	tgtgattgtt	ttttatagt	ttaatgtgtg	8700
tgagatgttg	gaagtattgg	ggattagtag	tagtttagat	gtttaaaaag	ataaggtgtt	8760
tttaatttgtg	tggattttatt	gaagttaagt	gggtgaataaa	gataattatt	tagataattt	8820
agattaaagt	aaaagtaaaa	ttatatattt	ttgtatatat	atattttatat	ttatttttata	8880
ttatagatat	atatatgtat	atatatatattg	gtttttgtaaa	taattgattt	aaagtgagga	8940
tttttttttgt	attttttttag	taggagtttt	aatattttttt	taattttttta	attatttttat	9000
a						9001

<210> 5

<211> 9001

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 5

tgtaaagtga	ttaggagatt	aggagaatgt	tgaaatttttt	gttggaaaaa	tgtaaagaaa	60
attttttattt	tgagtttagtt	gttttatagag	ttagtgtgtg	tgtgtgtgtg	tgtgtttgta	120
atataaaaatg	gatgtgaata	tatatatata	aatagatatg	gtttttgtttt	tatttttaatt	180
tgaattatttt	agataattgt	ttttattttat	tatttgattt	taatgggttt	atataaatta	240
ggatattttta	tttttttagg	tatttaggtt	gttggtgatt	tttagtggtt	ttaatattttt	300
gtatatgttg	gtattatgag	gagtagttat	gtgttttttg	gtttttttaat	tatttttgag	360
gttgattgag	gtttttttata	tatgtatat	tgtgtgtgat	aaagtttttat	tggtagagtg	420
gagttatttag	agttttttatt	aaaatttttgt	gggttttatga	gagatgggtt	tagaaattta	480
tatgggttttg	tgggggttttt	tgggtttttta	aaataaggta	ttaatatttta	agttttttaa	540
aatattttgta	gttttggggt	ttgaattttg	aaaaataagg	agtgaggggt	tgtgtatat	600
aattatagtg	gagattttttt	ttattttttta	atgtgatgga	gttttttttat	gaaatgaagt	660
tttaaggggt	atggatattgt	ggggattata	gttatttttga	ggtttaaaaag	aagaaatttg	720
aatatgatta	gtaaatatat	ttagtagaaa	agagttggat	ttttattgat	ttagttatag	780
gttatttggtt	ggtagtgtaa	tgggaggaaa	tattttatttt	atatatatat	tttatgattt	840
tgggggaatt	agaggaaatt	taataagaaa	atggttagaa	atattttaaaa	ttttttattta	900
aaagatttta	gtaaattaga	gtttttattag	attaaaaatt	attataaatg	taagagtatt	960
gttttttagtg	aaatgttgtg	gggttttgaga	aggagatttt	ttgttaaatt	tttgggataa	1020
aatgtgttat	tttaagtatta	gataatgagt	agaatgtaaa	tttaatttaat	ttttttttatt	1080
aataggttgt	tagtgtaattg	tgtataattt	agtataaga	ttgtaggatt	taatatagtt	1140
ggatgtatga	gttttagtta	atgtagattt	gttatatgag	gatgtgtttt	attttgagta	1200
ggtgtttgta	tgtgtggaat	ggggtaaagt	ggaataaaaag	gttaaaaagta	gaaatgttga	1260
tttaaaagttt	attatgaaga	aattttttttt	ttgtagttaa	attatttttta	aagtgggatg	1320
atattgggtga	agaaagattg	aaaaataatt	tttatgtgtg	tttttggttt	gtaagtttaa	1380
aatggggagg	agttgtagat	agggttttggg	ggtgggttagg	gtaaaggaga	gatataataag	1440
ttgtaaatat	atttgtagtt	tgtttttattt	attttggtttt	atattgaata	agtttttttaa	1500
ttttgtgaat	aaggataagg	agggagtgtt	ttaaagatat	tttatgttgg	tattgtaaat	1560
tattgattgt	aatgttaaat	aaatatatat	ttagagatga	taatatatta	tttatagtaa	1620
aataattgtt	tatgtagaaa	tttagaggag	attagtttgt	tttttttagt	tgattttatgt	1680
tgggggataa	aaggatttttt	aaaaattatt	ttgaatatgt	ttggatttttt	tttttttaatt	1740



tttttggaaa	ttaaatttgt	ttggaaatag	tgttataaag	agttgatggt	tttaaagggtg	1800
atTTTTTTTT	ttttatataa	ataagggttt	gtttttgtta	gttgagtgtg	gttttagggt	1860
ttttgttttt	agtttatata	tatttttttt	gtttgttttg	atTTtaatgg	tttaagatag	1920
ttttgagttt	attgggaaaa	gaaaatgatt	gttaaaaatt	atTTttgaaa	ttgggtattt	1980
ggtaatatTT	ttaatgtat	ggaaatttat	taagggtatat	tttatatata	attagtttaa	2040
ggttgttgat	tttatagggt	ttatggattt	aaatttgatt	gataataaag	taaataagag	2100
agttgaattt	aaagtgtggt	ttttttgggt	taggatgagt	ttaatatagt	gtataaggaa	2160
tttgaaagat	ttaggatatg	tgttttaatt	aatgttaagt	agaatggata	agtttttagt	2220
atTTtgaaaa	tgttgggtta	gggttttttt	tttattgtgt	gtttttgtgt	tggggattaa	2280
taagtattat	agagaatgtg	atTTgaggtg	atTTtttatt	tttgataaaa	tttagagtga	2340
attattaaat	agttgtttgt	ttaaagttaa	ggtaattttt	ttttgatggg	tttatttgtt	2400
ttttgatttt	taatttatta	gtttgttttt	ttagggtttt	gttttttttg	taattaaagt	2460
tttttttagat	tagtgtagta	tttatttgat	aggttgtttg	gaaaatttaa	gattggagag	2520
gtgatttgtt	gttggttttt	aaatttttta	gttttaagta	atgtgttttt	tttttatatg	2580
gggtggggga	ttggaaatgg	atgtagttag	atataaagag	tgggtgtttt	gttgattttt	2640
gtattttttt	ttttttgatt	atTTtatTTT	ttttttttta	gtttttgatt	tttagtttta	2700
tttttttatt	tttgggtttg	tattaaaagt	tggattgttt	tgggttgggt	aggagttaa	2760
tttttgggag	tttggttgtg	tagatttagt	gtgtatggtg	aggtagtagt	ttgggtttgt	2820
attgttgata	ggtgtaggta	ggatagtttt	tttattgtgg	tttgggggtg	tttgattggt	2880
gtggagttat	gttagttgta	tttggagaag	ggtttgggag	gaggtggagg	tggagagggt	2940
tggggagggt	tgtggtggag	tgatgttttg	gtattaggaa	gtttgttttt	ggttttaaga	3000
tgttaggtta	atagggaagt	gtggagttgt	agatttggtt	tgttgtttgt	ttgggtgttt	3060
ggagttgagt	tgtggttaagg	tttgggtttt	gtttgattgt	ttgaggggtg	tgtgtgtgtg	3120
tgttgtggag	ggtgtgttta	gagggttgtg	ttgtggttgt	agtgggttgt	gttgtttag	3180
gggatttaat	attatttatt	tgtttttgtt	atTTttgata	ttttttgttt	agggttgttg	3240
tgtggggggg	gggtgggtag	agtgtggttg	gtgttagttt	tttttattgg	aggggttttt	3300
gggggaggga	gggagagaag	aagggggttt	ttgtttattt	ttgtttgttt	ttggagtttg	3360
gaagtttgtt	ttttaaagat	gttttgagtg	gtgttttttt	gtttatatatt	tatgtttttg	3420
tttgtttgtt	gattttttgt	ttttggattt	ttttgtttga	gttttttggg	ggagatgggg	3480
gtagtttgggt	ttgagaattt	ggtgggggtt	gtgttttttt	gttttttttg	tagtggggaa	3540
atTTtgtgtt	tagagtgtga	tttggagtgg	gtagtggtgg	ttatgggggt	ttgggtgggt	3600
agtagttaag	gattagtaga	gtgttgtgtt	tttttgttta	tgaattgtat	gaaaggtttg	3660
ttttatttgg	agtattgagt	agtggggatt	aagtgttggg	ttgttttttt	atTTttttgt	3720
tattattttt	agttgttagt	tatggttttg	gttttgggtt	ttggtttagt	ttgggttgtg	3780
gatttttttt	agtatagggt	ggaggtgtat	attatttttg	atatttttag	tttggagggt	3840
gtaggtaagg	tgttgtgttg	ttttgtagat	atTTttgttt	agttgttttg	tgttatttgt	3900
ttttttttgt	tttaagggaag	ttagtttttt	tggggggagg	tgtggtggga	gtgggttgtt	3960
gtttgggttt	ttgtagaatt	tttgggaggt	ggaattttga	ttattttgta	tttttttagt	4020
ttttttttga	ttgggttgggt	ttttgggggt	ttaaagggtgt	gagtaatttt	gttgtttttt	4080
ttatttgtat	tttgggtttt	tttttgtttt	ttgggttata	aaaatttttag	tatttttgatt	4140
tgaggatttt	tagaggttgt	tgattttttgt	tttttgtttt	tttttgggtt	ttagtttttg	4200
aggagtttta	tttgttagga	aattgtttga	aattatttag	aaatgttttt	tgtgaagagg	4260
tatttttttt	tttttttttg	gaaagggttg	gtgaattttg	gtgtttaatt	gaatttttat	4320
atTTtttttt	agttttttta	aattgtatgg	aaatttgagt	tttttgtgag	ggggaggggg	4380
gtttgtaaat	tatgtgtgtg	tgtgtgtttt	aggagatttg	gtgtgtttgt	gtagagggtg	4440
ataaatatat	ttgaaagtat	aggttataaa	agtgaatgtg	ttgtttagt	gagataaata	4500
tgtaaataaa	atgtgtggtg	ttggggggagg	ggaggaaatg	gggtgtggat	atTTtatatt	4560
gtgtttgtat	atTTtatagg	tgtagtgttt	tttgtgggtt	ggagtgtttg	tgtgtatttt	4620
tttttgggtg	taggtagtgt	agttttttta	tgggtttttg	tgttgggtta	gttgggtgtt	4680
gtgtttagg	tgggtatgtt	gatgggaaag	tgtgtgtgtt	ttgttttttag	agaaagataa	4740
aagtttagtag	gggaagaatg	aggatgtggg	tgttgaggat	ttgtttaaga	agaagtggta	4800
aagggtgtag	tggatttatt	ttattagtta	gtagtttttag	gagttggagg	ttatttttta	4860
gaggaattgt	tatttggata	tgttttatatg	tgaagaaatt	gttgtgtgga	ttaatTTtat	4920
ggaagtttga	gtttgggtag	gagttagtat	ggagtttggg	agggatgggg	ggaggatgtt	4980
gtggagggtat	aggttaagta	gattaggaga	gaatgtggaa	ggtagtgttg	tttggggagg	5040
tgttgggtgg	gtgtagtttt	gtaaaggtag	aaggttttgt	gggtgggttg	ttgtgagatt	5100
atagtttttt	ttttgaggtt	gataggattg	ttgttttgggt	ttagggtttt	agagtgggtat	5160
tgggtttattg	ttttgttatt	ttgtgatttt	atgagttggg	ttgtatgggt	aattttttgt	5220
ataggatatt	gtgttttttg	tttgtagtgt	ttagagtaga	gttaataaaa	tttttatttag	5280
gttaagagtt	gtgaatagggt	tttaatttgt	gagtttttaa	taaggaaaat	ttgttagaga	5340
tatggaaagag	ttgggtttttt	ttgggaaatt	tttgtttttg	ttttgggttta	gttttttttt	5400
ttttgggttt	gtgtttttta	tatttttttt	atgggtgttt	tgggtatttta	ggtttttttt	5460
atataatttta	tttttttagtt	ttgtgatttt	tgggagtaaa	gttttaatat	ataattatta	5520

gttttttttag	aaggagaaaag	aaaaaaagaa	gaaagattttt	tttgttttggt	ttattttattt	5580
tttttttagga	gttgaattttt	ggaaattgaa	atttatatttt	tttttttttaa	attataatta	5640
tagtttttgta	aaaaggggttt	attttaatttt	tgtagtaaatt	ttgtattttta	tggattggta	5700
aaaatgagtt	taaataaata	atttaaatagt	aatgtttttgg	tttatgtttgg	ttggtggaag	5760
atttttaaatt	tgttaggattt	ttggaagtag	aaaatagaat	taagtaaatt	aagtgggtatt	5820
tagagggtttt	gttgttaaaa	aaaaaaaatt	aagtgtttttg	ggtagaaaaa	ataaagttttt	5880
tggtttagagt	agagtaaata	aaaagaagaa	aataatgata	aaaagaataa	agattaaaaat	5940
gttttttttaa	attagaggga	atgaagatat	tttttgggtg	gtattttgtgt	aagggtatgag	6000
gttatgtttgg	tggataaaaag	gttgggaaga	agttgaaaat	ggtttttagtt	taattgtttta	6060
gagtttagagt	tgggtttttgg	gtggtgtggt	tttgagtaag	gttagtttttt	tatttagttttt	6120
tttgtatatatt	aaggggaatgg	gtttttttatg	tattttttttt	gttttgagtaa	agtttagatg	6180
gttttagggta	gaaatggtaa	gtaattaaag	atagagtttta	tgggttttttt	gggattttttt	6240
gaaaatgtttt	ttttatttttg	tttgttatttt	tgtagttttta	tttttagtggt	ttgtagtgtgt	6300
gggtgttgggt	ttttttttgta	gttggtttttt	tttttaggggt	ggttgttttgt	tgagttaaagt	6360
gggagtgagg	tgtgttttttt	atagtagtttg	gggtgaaaga	ggaaggggga	taaaaaggaa	6420
attaagaatg	aaaggaaaaa	gagaaaaagt	ggattatatg	gttgggttttg	gtggagatgt	6480
gtaatgtgaa	atattattgg	tgttagtttg	gataatttttag	gttaggttttt	tttttaatat	6540
ataaaagtgtg	ttgttttgggg	tgataggggag	gtttgatgtg	gattgggatt	ggggttgttg	6600
ttgggttatt	ggatatgggt	ggaagtgtgt	tgggtttgggt	ggttgttttgt	aaagttaaatt	6660
gatttggttg	ggtttggtgt	gtggataggt	ttgtggtggg	tttagggtaa	agaagaggta	6720
gagtgaagaa	aggggggaatt	tttaaaatta	tttttttttgg	gttttttgag	tttaatatgt	6780
taagttttttg	gagttaatga	gttgatgaag	aggtggttttt	ttgttttttta	tttggttgtt	6840
ttgttaggtg	agaaagagtg	ttggtggttt	agttttttgtt	aagggagtat	gtattagggg	6900
gtgggggatg	atagtggagg	ttagggaagg	aagggaggaa	ttgtgtggga	gaaagagtga	6960
tttttttagtg	tttttttagt	ttttttttttt	tatttgtggg	tttgtggttt	tggaatggaa	7020
gtaagtttgt	aaggtgtttt	gggaagggtt	ggaaaagttt	gttggttttgt	gtttgttttta	7080
tattaagtgt	ttttggattt	ggagaaatgt	ttggttgagt	gattaaattg	tttgtagggt	7140
tttatgtgtt	tgggttgaggt	ttgtggtgta	gttttgagtt	ttagtttgta	ggttagagta	7200
gattagggtt	tttgtgtttg	gtggagattt	gggttagtaa	ttgaaagtgt	gttttggtat	7260
tttggtgtgt	aggggtggtgt	agtgaagtga	ggttaggggtg	tgtgagtgtg	ttagtgtgtg	7320
tgttggggga	aggtgggggt	tgggtttttga	tggaaagtttt	agtaatttgt	attgtggtat	7380
ttgtttgttt	ttttgttttta	attgtttttta	ggtttggttt	aagaattgtt	gggttaaattg	7440
gagaaagagg	gagtgttaatt	agtaggttga	gttatgtaag	aatgggttttg	ggttgtagtt	7500
taatgggttt	atgtagtttt	atgatgatat	gtatttaggt	tattttttata	ataattgggt	7560
tgttaaggggt	tttatatttg	ttttttttatt	tattaagagt	ttttttttttt	ttaatttttat	7620
gaatgttaatt	tttttgttat	tatagagtat	gttttttttta	tttaattttta	ttttgttttat	7680
gagtatgttg	tttagtatgg	tgttttttagt	agtgataggt	gtttttgggtt	ttagttttta	7740
tagtttgaat	aatttgaata	atttgagtag	tttggtgttg	aattttgttg	tgttgatgtt	7800
tgtttgtttt	tatgtgttgt	tgatttttttt	gtatgttttat	agggatatgt	gtaatttgag	7860
tttggttagt	ttgagattga	aagtaaagta	gtatttttagt	tttggttatg	ttagtgtgta	7920
gaatttggtt	tttaatttga	gtgtttgtta	gtatgtagtg	gattgggtttg	tgtgagttgt	7980
atttatagtg	ttgggattttt	aggatttttgt	tggatgggggt	aattttgtttt	ttgaaagatt	8040
gggaatttatg	ttagaagggtt	gtgggtatta	aagaaaggga	gagaaagaga	agttatatag	8100
agaaaaggaa	attattgaat	taaagagaga	gttttttttga	ttttaaagggt	atgttttttag	8160
tgtttgatatt	tttttatttat	aagtatttttt	aatagtttga	aggatatata	tataaataaaa	8220
tgtttgattg	gatattgatat	tttaatatata	ttataagttt	gttattttttt	aagttttagta	8280
ttgttaatat	ttaaatgatt	gaaaggatgt	atatatatattg	aaatgttaaaa	tttaatttttat	8340
aaaagtagtt	gttagtaata	ttataaatagt	gttttttaaaag	gttaggttttt	aaaataaagt	8400
atgttatata	gaagtgatta	ggatttttttg	tttgtgagta	agggagtgtga	tatatataaat	8460
gttatattgt	atgttttttaa	tatatattata	ttattataaaa	aaatgtgtga	atattagttt	8520
tagaatagtt	tttttgggtgg	atgtaaatgat	gttttttgaaa	ttgttatgta	taattttattt	8580
tgtgtataaat	attttgtata	atattatttgt	tttattttttt	agtaaataatg	aaataaaatgt	8640
gtttttatttt	atgggagtaa	aatatatattgt	atataaatttg	gtttggatttt	ttttttttttt	8700
tttttgttat	taatttgggtt	aggatatattt	agttatttgtt	ttttaaataa	attagtttttt	8760
tttggttgtt	tagttaaata	tataaggtag	tagttttttat	ttaaattttgg	tagaaataaaa	8820
tgatagttat	ttattagaaa	ttaaaaagaa	aaaaaaagggt	attttttgggg	gggaaaagggt	8880
ttataaaaatt	taattttgtt	tttttaatttt	tttttttggtt	taaattttaga	ggatttttatt	8940
atggtttagta	aataatatga	aaaagaaaaa	agaagaaaga	aattttagtaa	gtttatttagt	9000
t						9001

<210> 6  
<211> 22  
<212> DNA

<213> Artificial Sequence

<220>  
<223> PRIMER

<400> 6

gtaggggagg gaagtagatg tt 22

<210> 7  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PRIMER

<400> 7

ttctaatacct cctttccaca ataa 24

<210> 8  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PROBE

<400> 8

agtcggagtc gggagagcga 20

<210> 9  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PROBE

<400> 9

agttggagtt gggagagtga aaggaga 27

<210> 10  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PRIMER CONTROL

<400> 10

tggtgatgga ggaggtttag taagt 25

<210> 11  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>

<223> PRIMER CONTROL

<400> 11

aaccaataaa acctactcct cccttaa

27

<210> 12

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> PROBE CONTROL

<400> 12

accaccaccc aacacacaat aacaaacaca

30

<210> 13

<211> 408

<212> DNA

<213> Homo Sapiens

<400> 13

tcctcaactc	tgcaggcctg	aaagaaggtc	acacacgcac	gctcacaccc	acactccaca	60
cgctcgtcc	caaacaaccc	catgaacatt	gtcctttgtt	ccgtctcttg	ggccactttc	120
cctgtogett	cctcccagcc	cgctctgatt	tgctcccaa	aagtacgttt	ctgtctcccc	180
gctgccttg	cgctccccct	ttgatttatt	agggtgccc	ggttggcgca	gattgctttt	240
tcttctcttc	catcccatcc	tcccttcttg	tcctcctttc	cacagtggga	gtccgtgctc	300
ctgctcctcg	gttggctcct	aagtgcccc	ccaggcccc	tctcctttcg	ctctcccggc	360
tccggctccc	gactcttcgg	cccgtggca	tctgcttccc	tcccctgc		408

<210> 14

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 14

aacatctact tccctcccct ac

22

<210> 15

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 15

gtagtagag attttattaa attttattgt at

32

<210> 16

<211> 14

<212> DNA

<213> Artificial Sequence

<220>



<223> chemically treated genomic DNA (Homo sapiens)

<400> 16

ttcggttgcg cggt

14

<210> 17

<211> 16

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 17

tttggttggtg tggttg

16

<210> 18

<211> 144

<212> DNA

<213> Homo Sapiens

<400> 18

ttctggtcct cctttccaca gtgggagtcg gtgctcctgc tcctcggttg gctcctaagt  
gccccgccag gtcccctctc ctttcgctct cccggctccg gctcccgaact cttcggccccg  
ctggcatctg ctteccctccc ctgc

60  
120  
144

<210> 19

<211> 162

<212> DNA

<213> Homo Sapiens

<400> 19

tggcatctgc ttccctcccc tgcctcgttt ctgcctgcgcc ctgctcgctc cccccggcgc  
tcgcccgggc gctgtgctcg ctccctggatc gccagccgcg cagcgggctc gccggcgccc  
gcgcgccact gtgcagtgga gtttggtgga atctctgctg ac

60  
120  
162

<210> 20

<211> 2235

<212> DNA

<213> Homo Sapiens

<400> 20

tgggagtcgc	tgctcctgct	cctcgggttg	ctcctaagt	ccccgccagg	tcccctctcc	60
tttcgctctc	ccggctccgg	ctcccgaact	ttcggccccg	tggcatctgc	ttccctcccc	120
tgcctcgttt	ctcgtcgccc	ctgctcgctc	cccccgggcg	tcgcccgggc	gctgtgctcg	180
ctcctggatc	gccagccgcg	cagccgggct	cggccggccc	cccgcgcgcc	actgtgcagt	240
ggagtttggt	ggaatctctg	ctgacgtcac	gtcactcccc	acacggagta	ggagcagagg	300
gaagagagag	ggatgagagg	gagggagagg	agagagagtg	cgagaccgag	cgagaaagct	360
ggagaggagc	agaaagaaac	tgccagtggc	ggctagattt	cggaggcccc	agtgcacccc	420
tggactcctt	cggaaacttg	caccctcagg	agccctgcag	tcctctcagg	cccggctttc	480
gggcgccttg	cgtgcagccg	gaggctcggc	tcgctggaaa	tcgccccggg	aagcagtggg	540
acgcggagac	agcagctctc	tcccggtagc	cgataacggg	gaatggagac	caactgccgc	600
aaactggtgt	cggcgtgtgt	gcaattaggc	gtgcagccgg	cggccggttg	atgtctcttc	660
tccaaagact	ccgaaatcaa	aaaggctcag	ttcacggact	ctcctgagag	ccgaaaagag	720
gcagccagca	gcaagttctt	cccgcggcag	catcctggcg	ccaatgagaa	agataaaaag	780
cageagggga	agaatgagga	cgtgggccc	gaggacccgt	ctaagaagaa	gcggcaaagg	840
cggcagcggg	ctcactttac	cagccagcag	ctccaggagc	tggaggccac	tttccagagg	900
aaccgctacc	cggacatgtc	cacacgcgaa	gaaatcgctg	tgtggacca	ccttacggaa	960

gcccagagtc	gggtttgggt	caagaatcgt	cgggccaaat	ggagaaagag	ggagcgcaac	1020
cagcaggccg	agctatgcaa	gaatggcttc	gggccgcagt	tcaatgggct	catgcagccc	1080
tacgacgaca	tgtaccagag	ctattcctac	aacaactggg	ccgccaaggg	ccttacatcc	1140
gcctccctat	ccaccaagag	cttccccttc	ttcaactcta	tgaacgtcaa	ccccctgtca	1200
tcacagagca	tgttttcccc	acccaactct	atctcgtcca	tgagcatgtc	gtccagcatg	1260
gtgccctcag	cagtgcacag	cgtcccgggc	tccagtctca	acagcctgaa	taacttgaac	1320
aacctgagta	gcccgtcgct	gaattccgcg	gtgccgacgc	ctgcctgtcc	ttacgcgccg	1380
ccgactcctc	cgtatgttta	tagggacacg	tgtaactcga	gcctggccag	cctgagactg	1440
aaagcaaagc	agcactccag	cttcgggtac	gccagcgtgc	agaacccggc	ctccaacctg	1500
agtgccttgc	agtatgcagt	ggaccggccc	gtgtgagccg	caccacacgc	gccgggatcc	1560
taggaccttg	ccggatgggg	caactccgcc	cttgaaagac	tgggaattat	gctagaaggt	1620
cgtgggcact	aaagaaaggg	agagaaagag	aagctatata	gagaaaagga	aaccactgaa	1680
tcaaagagag	agctcctttg	atttcaaagg	gatgtcctca	gtgtctgaca	tctttcacta	1740
caagtatttc	taacagttgc	aaggacacat	acacaaacaa	atgtttgact	ggatatgaca	1800
ttttaacatt	actataagct	tgttatTTTT	taagttagtc	attgttaaca	tttaaataac	1860
tgaaaggatg	tatatatatc	gaaatgtcaa	attaatttta	taaaagcagt	tgtagtaaat	1920
atcacaacag	tgttttttaa	ggttaggctt	taaaataaag	catgtttata	agaagcgatt	1980
aggatttttc	gcttgcgagc	aaggggagtgt	atatactaaa	tgccacactg	tatgtttcta	2040
acatattatt	attattataa	aaaatgtgtg	aatatcagtt	ttagaatagt	ttctctgggt	2100
gatgcaatga	tgtttctgaa	actgctatgt	acaacctacc	ctgtgtataa	catttcgtac	2160
aatattattg	ttttactttt	cagcaaatat	gaaacaaatg	tgttttattt	catgggagta	2220
aaatatactg	catac					2235

<210> 21

<211>

<212> protein

<213> polypeptide Sequence

<220>

<223> amino acid sequence (Homo sapiens)

<400> 450

METNCRKLVSAQVQLGVQPAAVECLFSKDSEIKKVEFTDSPESRKEAASSKFFP  
 RQHPGANEKDKSQGKNEVDGAEDPSKKKRQRRQRTHTFSQQQLQELEATFQRNR  
 YPDMSTREEIAVWTNLTEARVRVWFKNRRRAKWRKRERNQQAELCKNGFGPQFNG  
 LMQPYDDMYPGYSYNNWAAKGLTSASLSTKSFPFFNSMNVNPLSSQSMFSPNS  
 ISSMSMSSSMVPSAVTGVPGLSLNLSLNNLNLSSPSLNSAVPTPACPYAPPTPP  
 YVYRDTCNSSLASRLKAKQHSFSGYASVQNPASNLSACQYAVDRPV

450

<210> 22

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 22

gtaggggagg gaagtagatg t

21

<210> 23

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 23

tcctcaactc taaaaaccta aaa

23

<210> 24  
<211> 16  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> chemically treated genomic DNA (Homo sapiens)

<400> 24

agtcgggaga gcgaaa

16

<210> 25  
<211> 16  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> chemically treated genomic DNA (Homo sapiens)

<400> 25

agttgggaga gtgaaa

16

<210> 26  
<211> 17  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> chemically treated genomic DNA (Homo sapiens)

<400> 26

aagagtcggg agtcgga

17

<210> 27  
<211> 17  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> chemically treated genomic DNA (Homo sapiens)

<400> 27

aagagttggg agttgga

17

<210> 28  
<211> 16  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> chemically treated genomic DNA (Homo sapiens)

<400> 28

ggtcgaagag tcggga

16

---

<210> 29  
<211> 16

<212> DNA  
<213> Artificial Sequence

<220>  
<223> chemically treated genomic DNA (Homo sapiens)

<400> 29

ggttgaagag ttggga

16

<210> 30  
<211> 16  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> chemically treated genomic DNA (Homo sapiens)

<400> 30

atggttagcgg gtcgaa

16

<210> 31  
<211> 16  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> chemically treated genomic DNA (Homo sapiens)

<400> 31

tagtggggttg aagagt

16

<210> 32  
<211> 24  
<212> DNA  
<213> Homo Sapiens

<400> 32

gaaaggcaga gtcataacag gaag

24

<210> 33  
<211> 26  
<212> DNA  
<213> Homo Sapiens

<400> 33

taaggataga gtgatttcca agaaag

26

<210> 34  
<211> 24  
<212> DNA  
<213> Homo Sapiens

<400> 34

cttatctgac aacaagcgag tatg

24

<210> 35  
<211> 25



<212> DNA

<213> Homo Sapiens

<400> 35

caattattgt attgtagcat cggag